

University of Ottawa  
Department of Mathematics and Statistics

MAT 1302E : Mathematical Methods II  
Professor: Catalin Rada

Second Midterm Exam – Version A

November 21, 2013

Surname \_\_\_\_\_ First Name \_\_\_\_\_

Student # \_\_\_\_\_ DGD (Monday/Thursday) \_\_\_\_\_

**Instructions:**

- (a) You have 80 minutes to complete this exam.
- (b) The number of points available for each question is indicated in square brackets.
- (c) All work to be considered for grading should be written in the space provided. The reverse side of pages is for scrap work. If you find that you need extra space in order to answer a particular question, you should continue on the reverse side of the page and indicate this **clearly**. Otherwise, the work written on the reverse side of pages will not be considered for marks.
- (d) Write your student number at the top of each page in the space provided.
- (e) No notes, books, scrap paper, calculators or other electronic devices are allowed.
- (f) You are strongly recommended to write in **pen**, not pencil.
- (g) You may use the last page of the exam as scrap paper.

Good luck!

Please do not write in the table below.

Question	1	2	3	4	5	6	Total
Maximum	3	3	4	5	5	3	23
Grade							

1. [3 points] Find the inverse of the matrix

$$A = \begin{bmatrix} 1 & 3 & -2 \\ 1 & 1 & -2 \\ 2 & 1 & -3 \end{bmatrix}.$$

**Solution:**

$$\begin{aligned}
[A|I] &= \left[ \begin{array}{ccc|ccc} 1 & 3 & -2 & 1 & 0 & 0 \\ 1 & 1 & -2 & 0 & 1 & 0 \\ 2 & 1 & -3 & 0 & 0 & 1 \end{array} \right] \xrightarrow{\substack{R_2 \rightarrow R_2 - R_1 \\ R_3 \rightarrow R_3 - 2R_1}} \left[ \begin{array}{ccc|ccc} 1 & 3 & -2 & 1 & 0 & 0 \\ 0 & -2 & 0 & -1 & 1 & 0 \\ 0 & -5 & 1 & -2 & 0 & 1 \end{array} \right] \\
&\xrightarrow{R_3 \rightarrow R_3 - \frac{5}{2}R_2} \left[ \begin{array}{ccc|ccc} 1 & 3 & -2 & 1 & 0 & 0 \\ 0 & -2 & 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{5}{2} & 1 \end{array} \right] \xrightarrow{R_1 \rightarrow R_1 + 2R_3} \left[ \begin{array}{ccc|ccc} 1 & 3 & 0 & 2 & -5 & 2 \\ 0 & -2 & 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{5}{2} & 1 \end{array} \right] \\
&\xrightarrow{R_2 \rightarrow -\frac{1}{2}R_2} \left[ \begin{array}{ccc|ccc} 1 & 3 & 0 & 2 & -5 & 2 \\ 0 & 1 & 0 & \frac{1}{2} & -\frac{1}{2} & 0 \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{5}{2} & 1 \end{array} \right] \xrightarrow{R_1 \rightarrow R_1 - 3R_2} \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & -\frac{7}{2} & 2 \\ 0 & 1 & 0 & \frac{1}{2} & -\frac{1}{2} & 0 \\ 0 & 0 & 1 & \frac{1}{2} & -\frac{5}{2} & 1 \end{array} \right]
\end{aligned}$$

$$\text{Therefore } A^{-1} = \begin{bmatrix} \frac{1}{2} & -\frac{7}{2} & 2 \\ \frac{1}{2} & -\frac{1}{2} & 0 \\ \frac{1}{2} & -\frac{5}{2} & 1 \end{bmatrix}$$

2. [3 points] Let  $A = \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$ ,  $B = \begin{bmatrix} 0 & 1 \\ 2 & 2 \end{bmatrix}$  and  $C = \begin{bmatrix} 1 & -1 & 3 \\ 2 & 0 & 1 \end{bmatrix}$ . Solve the following matrix equation for  $X$ :

$$A(X^T + C) = BC$$

**Solution:**

$$\begin{aligned} A(X^T + C) = BC &\Rightarrow A^{-1}A(X^T + C) = A^{-1}BC \Rightarrow X^T + C = A^{-1}BC \\ &\Rightarrow X^T = A^{-1}BC - C \Rightarrow X = C^T B^T (A^{-1})^T - C^T \end{aligned}$$

Now we have

$$A^{-1} = \frac{1}{1} \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 1 & 1 \end{bmatrix} \Rightarrow (A^{-1})^T = \begin{bmatrix} 3 & 1 \\ 2 & 1 \end{bmatrix}$$

and therefore

$$C^T B^T (A^{-1})^T = \begin{bmatrix} 1 & 2 \\ -1 & 0 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} 0 & 2 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} 18 & 8 \\ -4 & -2 \\ 19 & 9 \end{bmatrix}$$

and

$$X = \begin{bmatrix} 18 & 8 \\ -4 & -2 \\ 19 & 9 \end{bmatrix} - \begin{bmatrix} 1 & 2 \\ -1 & 0 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} 17 & 6 \\ -3 & -2 \\ 16 & 8 \end{bmatrix}$$

3. Let  $A = \begin{bmatrix} 1 & 2 & 4 & 0 & 3 \\ 0 & 0 & -2 & 1 & -1 \\ 0 & 0 & 2 & -1 & 1 \end{bmatrix}$ .

(a) [3 points] Find a basis for  $\text{Nul } A$ . **Solution:** The reduced echelon form of  $A$  is

as follows:

$$\begin{aligned} \begin{bmatrix} 1 & 2 & 4 & 0 & 3 \\ 0 & 0 & -2 & 1 & -1 \\ 0 & 0 & 2 & -1 & 1 \end{bmatrix} &\xrightarrow{R_3 \rightarrow R_3 + R_2} \begin{bmatrix} 1 & 2 & 4 & 0 & 3 \\ 0 & 0 & -2 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_2 \rightarrow -\frac{1}{2}R_2} \begin{bmatrix} 1 & 2 & 4 & 0 & 3 \\ 0 & 0 & 1 & -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \\ &\xrightarrow{R_1 \rightarrow R_1 - 4R_2} \begin{bmatrix} 1 & 2 & 0 & 2 & 1 \\ 0 & 0 & 1 & -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \end{aligned}$$

In the corresponding homogeneous linear system, the variables  $x_1$  and  $x_3$  are basic, and  $x_2, x_4, x_5$  are free. The general solution is

$$\begin{cases} x_1 = -2x_2 - 2x_4 - x_5 \\ x_3 = \frac{1}{2}x_4 - \frac{1}{2}x_5 \end{cases}$$

and its vector parametric form is

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -2 \\ 0 \\ \frac{1}{2} \\ 1 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -1 \\ 0 \\ -\frac{1}{2} \\ 0 \\ 1 \end{bmatrix}.$$

Therefore a basis for  $\text{Nul } A$  is

$$\left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ \frac{1}{2} \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ -\frac{1}{2} \\ 0 \\ 1 \end{bmatrix} \right\}.$$

(b) [1 point] What is  $\text{rank } A$ ? Justify your answer.

**Solution:** By the rank-nullity theorem we know that

$$\text{rank } A + \dim \text{Nul } A = 5 \Rightarrow \text{rank } A = 5 - 3 = 2.$$

4. An economy consists of two sectors: the Construction sector and the Service sector. We know that:

- For each unit of output, Construction requires  $\frac{1}{3}$  units from itself and  $\frac{1}{2}$  units from Service.
- For each unit of output, Service requires  $\frac{1}{6}$  units from Construction and  $\frac{3}{4}$  units from itself.

- (a) [1 point] What is the consumption matrix  $C$  for this economy?
- (b) [2 points] Find  $I - C$  and  $(I - C)^{-1}$ .
- (c) [1 point] Determine what intermediate demands are created if Construction wants to produce 12 units and Service wants to produce 24 units.
- (d) [1 point] Find the production levels that will satisfy the final demand of 5 units from Construction and 10 units from Service.

**Solution:** (a) The consumption matrix of this economy is:

$$C = \begin{bmatrix} \frac{1}{3} & \frac{1}{6} \\ \frac{1}{2} & \frac{3}{4} \end{bmatrix}.$$

(b)

$$I - C = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} \frac{1}{3} & \frac{1}{6} \\ \frac{1}{2} & \frac{3}{4} \end{bmatrix} = \begin{bmatrix} \frac{2}{3} & -\frac{1}{6} \\ -\frac{1}{2} & \frac{1}{4} \end{bmatrix}.$$

$$(I - C)^{-1} = \frac{1}{\frac{2}{3} - \frac{1}{12}} \begin{bmatrix} \frac{1}{4} & \frac{1}{6} \\ \frac{1}{2} & \frac{2}{3} \end{bmatrix} = 12 \begin{bmatrix} \frac{1}{4} & \frac{1}{6} \\ \frac{1}{2} & \frac{2}{3} \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 6 & 8 \end{bmatrix}.$$

(c)

$$12 \begin{bmatrix} \frac{1}{3} \\ \frac{1}{2} \end{bmatrix} + 24 \begin{bmatrix} \frac{1}{6} \\ \frac{3}{4} \end{bmatrix} = \begin{bmatrix} 8 \\ 24 \end{bmatrix}$$

(d) Let  $x_1$  and  $x_2$  be the outputs of Construction and Service respectively. Let  $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$  be the production vector. Note that  $\mathbf{d} = \begin{bmatrix} 5 \\ 10 \end{bmatrix}$ .

$$\mathbf{x} = (\mathbf{I} - \mathbf{C})^{-1} \mathbf{d} = \begin{bmatrix} 3 & 2 \\ 6 & 8 \end{bmatrix} \begin{bmatrix} 5 \\ 10 \end{bmatrix} = \begin{bmatrix} 35 \\ 110 \end{bmatrix}.$$

5. Consider the matrix

$$A = \begin{bmatrix} -1 & -1 & 0 & 2 \\ 1 & 3 & 1 & -2 \\ 2 & 1 & 0 & -3 \\ 0 & 4 & -1 & 5 \end{bmatrix}.$$

(a) [3 points] Calculate  $\det A$  by using the cofactor expansion method.

**Solution:** We use cofactor expansion along the third column:

$$\begin{aligned} \det A &= 0 \begin{vmatrix} 1 & 3 & -2 \\ 2 & 1 & -3 \\ 0 & 4 & 5 \end{vmatrix} - 1 \begin{vmatrix} -1 & -1 & 2 \\ 2 & 1 & -3 \\ 0 & 4 & 5 \end{vmatrix} + 0 \begin{vmatrix} -1 & -1 & 2 \\ 1 & 3 & -2 \\ 0 & 4 & 5 \end{vmatrix} - (-1) \begin{vmatrix} -1 & -1 & 2 \\ 1 & 3 & -2 \\ 2 & 1 & -3 \end{vmatrix} \\ &= - \begin{vmatrix} -1 & -1 & 2 \\ 2 & 1 & -3 \\ 0 & 4 & 5 \end{vmatrix} + \begin{vmatrix} -1 & -1 & 2 \\ 1 & 3 & -2 \\ 2 & 1 & -3 \end{vmatrix}. \end{aligned}$$

Now we have

$$\begin{vmatrix} -1 & -1 & 2 \\ 2 & 1 & -3 \\ 0 & 4 & 5 \end{vmatrix} = 0 \begin{vmatrix} -1 & 2 \\ 1 & -3 \end{vmatrix} - 4 \begin{vmatrix} -1 & 2 \\ 2 & -3 \end{vmatrix} + 5 \begin{vmatrix} -1 & -1 \\ 2 & 1 \end{vmatrix} = (-4)(-1) + (5)(1) = 9 \text{ (last row expansion)}$$

and

$$\begin{vmatrix} -1 & -1 & 2 \\ 1 & 3 & -2 \\ 2 & 1 & -3 \end{vmatrix} = (-1) \begin{vmatrix} 3 & -2 \\ 1 & -3 \end{vmatrix} - (-1) \begin{vmatrix} 1 & -2 \\ 2 & -3 \end{vmatrix} + 2 \begin{vmatrix} 1 & 3 \\ 2 & 1 \end{vmatrix} = (-1)(-7) + (1)(1) + (2)(-5) = -2. \text{ (first row)}$$

Therefore  $\det A = (-9) + (-2) = -11$ .

Student # \_\_\_\_\_

MAT 1302E Second Midterm Exam

(b) [2 points] This part is independent of part (a).

Let  $P, Q, R$  be  $3 \times 3$  matrices such that  $\det(P) = -\frac{1}{2}$ ,  $\det(Q) = 5$ , and  $\det(R) = -10$ . Compute  $\det(-\frac{1}{2}P^T Q^2 R^{-1} P^{-2})$ .

**Solution:**

$$\det(-\frac{1}{2}P^T Q^2 R^{-1} P^{-2}) = (-\frac{1}{2})^3 (-\frac{1}{2})(5^2)(-\frac{1}{10})(-2)^2 = -\frac{10}{16} = -\frac{5}{8}.$$

6. Determine if the following sets are subspaces of  $\mathbb{R}^3$  or not. Justify your answer.

(a) [1 point]

$$U = \left\{ \begin{bmatrix} a^2 + b^2 \\ b + c \\ 2c \end{bmatrix} \mid a, b, c \in \mathbb{R} \right\}$$

**Solution:** The top entry of every vector in  $U$  is  $a^2 + b^2 \geq 0$ . Therefore  $U$  is not a subspace because it is not closed under scaling:

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \text{ is in } U \text{ for } a = 1, b = c = 0$$

but

$$(-1) \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 0 \end{bmatrix} \text{ is not in } U.$$

(b) [1 point]  $V = \left\{ \begin{bmatrix} a - 2b \\ -b + c \\ -2c \end{bmatrix} \mid a, b, c \in \mathbb{R} \right\}$

**Solution:**  $V$  is a subspace because every vector in  $V$  can be written as

$$\begin{bmatrix} a - 2b \\ -b + c \\ -2c \end{bmatrix} = a \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + b \begin{bmatrix} -2 \\ -1 \\ 0 \end{bmatrix} + c \begin{bmatrix} 0 \\ 1 \\ -2 \end{bmatrix}$$

and therefore

$$V = \text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -2 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -2 \end{bmatrix} \right\}$$



(c) [1 point]

$$W = \left\{ \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \mid x_1, x_2, x_3 \in \mathbb{R} \text{ and } \begin{bmatrix} 1 & 0 & -3 \\ 2 & 5 & -2 \\ 5 & 10 & -7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right\}$$

**Solution:** Note that  $W = \text{Nul } A$  where  $A$  is the given  $3 \times 3$  matrix. Therefore  $W$  is a subspace.