

## OXYGEN AND CARBON ISOTOPES REFERENCE LIST

\* Lowenstam is the only reference that has  $\delta^{18}\text{O}$  but no  $\delta^{13}\text{C}$  Values.

Adlis, D.S.; Grossman, E.L.; Yancey, T.E.; McLerran, D.R. (1988): Isotope stratigraphy and paleodepth changes of Pennsylvanian cyclical sedimentary deposits. *Palaios*, 3, 487-506.

Brand, U. and Legrand-Blain, M., 1993. Paleocology and biogeochemistry of brachiopods from the Devonian-Carboniferous boundary interval of the Griotte Formation, La Serre, Montagne Noire, France. In: M. StreeL, G. Sevastopulo G. and E. Paproth (Editors),

Brand, U., 1989a. Aragonite-calcite transformation based on Pennsylvanian molluscs. *Geol. Soc. Amer. Bull.*, 101, 377-390.

Brand, U., 1989b. Biogeochemistry of late Paleozoic North American brachiopods and secular variation of seawater composition. *Biogeochemistry*, 7: 159-193.

Brand, U.; Veizer, J. (1981): Chemical diagenesis of a multicomponent carbonate system - 2: Stable isotopes. *J. Sed. Petr.*, 50, 987-997.

Carden, G. A. F., 1994, Stable isotopic changes across the Ordovician-Silurian boundary Ph.D.dissert. , The University of Liverpool, 284 p.

Carpenter, S. J. and Lohmann, K. C., 1995.  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  values of modern brachiopods. *Geochim. Cosmochim. Acta*, 59: 3749-3764.

Compston, W., 1960. The carbon isotopic composition of certain marine invertebrates and coals from the Australian Permian. *Geochim. Cosmochim. Acta*, 18: 1-22.

Delaney, M. L., Popp, B. N., Lepzelter, C. G. and Anderson, T. F., 1989. Lithium-to-calcium ratios in modern Cenozoic, and Paleozoic articulate brachiopod shells. *Paleoceanography* 4: 681-691

Douglas, R. G. and Savin, S. M., 1971. Isotopic analysis of planktonic foraminifera from the Cenozoic of northwest Pacific, Leg 6. Initial Rep. Deep See Drill. Proj. 6, 1123-1127.

Douglas, R. G. and Savin, S. M., 1973. Oxygen and Carbon isotope analyses of Cretaceous and Tertiary foraminifera from Central North Pacific. Initial Rep. Deep Drill. Proj. 17, 591-605.

Grossman, E.L., Mii, H-S. and Yancey, T.E., 1993. Stable isotopes in late Pennsylvanian brachiopods from the United States: Implications for Carboniferous paleoceanography. *Geol. Soc. Am. Bulletin*, 105: 1284-1392

Grossman, E.L., Zhang, C. and Yancey, T.E., 1991. Stable-isotope stratigraphy of brachiopods from Pennsylvanian shales in Texas. *Geol. Soc. Am. Bulletin*, 103: 953-965.

James, N. P., Bone, Y., Kyser, T. K., 1997. Brachiopod  $\delta^{18}\text{O}$  values do reflect ambient oceanography: Lacedpede Shelf, southern Australia. *Geology*, 25: 551-554

Jones, C. E., 1992. Strontium isotopes in Jurassic and Early Cretaceous seawater. Unpublished PhD thesis. University of Oxford.

Lavoie, D., 1993. Early Devonian marine isotopic signatures: Brachiopods from the upper Gaspé limestones, Gaspé Peninsula, Quebec, Canada. *J. Sediment. Petrol.*, 63: 620-627

Lowenstam, H.A., 1961. Mineralogy, O-18/O-16 ratios, and strontium and magnesium contents of recent and fossil brachiopods and their bearing on the history of oceans. *J. Geol.*, 69: 241-260

Middleton, P. D., Marshall, J. D. and Brechley, P. J., 1991. Evidence for isotopic changes associated with Late Ordovician glaciation, from brachiopods and marine cements of central Sweden. In Barnes, C. R. and Williams, S. H. (eds.) *Advances in Ordovician Geology*, Geol. Surv. Can. Pap., 90-9: 313-323.

Mii, H. S., (1996). Late Paleozoic environments: Carbon and Oxygen isotope records and elemental concentrations of brachiopod shells. PhD thesis, Texas A and M University, 191 pp., College station

Miller, K. G., Fairbanks, R. G. and Mountain, G. S., 1987. Tertiary oxygen isotope synthesis, sea level history, and continental margin erosion. *Paleoceanography*, 2: 1-19.

Mii, H.-S.; Grossman, E.L.; Yancey, T.E., 1997. Stable carbon and oxygen isotope shifts in Permian seas of West Spitsbergen-Global change or diagenetic artefact. - *Geology* 25, 227-230.

Miller, K.G., Fairbanks, R.G. and Mountain, G.S., 1987. Tertiary oxygen isotope synthesis, sea level history, and continental margin erosion, *Paleoceanography*, 2: 1-19.

Popp, B. N., Anderson, T. F. and Sandberg, P. A., 1986b. Textural, elemental and isotopic variations among constituents in Middle Devonian limestones, North America. *J. Sediment. Petrol.*, 56: 715-727.

Popp, B.N., Anderson, T.F. and Sandberg, P.A., 1986a. Brachiopods as indicators of original isotopic compositions in some Paleozoic limestones. *Geol. Soc. Am. Bulletin*, 97: 1262-1269.

Popp, N.B., Podosek, F.A., Brannon, J.C., Anderson, T.F. and Pier, J. 1986c.  $^{87}\text{Sr}/^{86}\text{Sr}$  in Permo-Carboniferous seawater from the analyses of well preserved brachiopod shells. *Geochim. Cosmochim. Acta*, 50: 1321-1328.

Qing, H. and Veizer, J., 1994. Oxygen and carbon isotope composition of Ordovician brachiopods: Implications for coeval seawater. *Geochim. Cosmochim. Acta*, 58: 1501- 1509.

Qing, H, Barnes C. R., Buhl, D., and Veizer, J., 1998. The Strontium isotopic composition of Ordovician and Silurian brachiopods and conodonts: relationships to geological events and implications for coeval seawater. *Geochim. et Cosmochim. Acta* 62: 1721-1734.

Rao, C. P., 1996. Oxygen and carbon isotope composition of skeletons from temperate shelf carbonates, eastern Tasmania, Australia. *Carbonates and Evaporites*, 11: 169-181

Rush, P. F. and Chafetz, H. S., 1990. Fabric-retentive, non-luminescent brachiopods as indicators of original  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  composition: A test. *J. Sed. Petrol.*, 60: 968-981.

Samtleben, C., Munneke, A., Bickert, T. and Pätzold, J., 1996, The Silurian of Gotland (Sweden): Facies interpretation based on stable isotopes in brachiopod shells: *Geologische Rundschau*, v. 85, p. 278-292.

Veizer, J. and Hoefs, J. (1976): The nature of  $\text{O}^{18}/\text{O}^{16}$  and  $\text{C}^{13}/\text{C}^{12}$  secular trends in sedimentary carbonate rocks. *Geochimica et Cosmochimica Acta* 40; 1387-1395.

Veizer, J., Fritz, P. and Jones, B., 1986, Geochemistry of brachiopods: oxygen and carbon isotopic records of Paleozoic oceans: *Geochimica et Cosmochimica Acta*, v. 50, p. 1679-1996.

Wadleigh M.A., Veizer J. (1992)  $^{18}\text{O}/^{16}\text{O}$  and  $^{13}\text{C}/^{12}\text{C}$  in Lower Paleozoic brachiopods: isotopic composition of sea water. *Geochim. et Cosmochim. Acta* 56, 431-487

Wenzel, B and Joachimski, M. M., 1996, Carbon and oxygen isotopic composition of Silurian brachiopods (Gotland/Sweden)- palaeoceanographic implications: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 122, p. 143-166.

### Sr ISOTOPES REFERENCE LIST

Banner, J.L. and Kaufman, J. (1994) The isotopic record of ocean chemistry and diagenesis preserved in non-luminescent brachiopods from Mississippian carbonate rocks, Illinois and Missouri. *Geol. Soc. Am. Bull.* 106, 1074-1082.

Barrera E., Baldauf J. and Lohmann K.C. (1993) 15. Strontium isotope and benthic foraminifer stable isotope results from oligocene sediment at site 803. *Proc. of the ODP, Sci. Res.* 130, 269-279.

Bertram, C. J., Elderfield, H., Aldridge, R. J., and Morris, S. C., 1992,  $^{87}\text{Sr}/^{86}\text{Sr}$ ,  $^{143}\text{Nd}/^{144}\text{Nd}$  and REEs in Silurian phosphatic fossils: *Earth and Planetary Science Letters*, v. 113, p. 239-249.

Brand, U., 1991. Strontium isotope diagenesis of biogenic aragonite and low-Mg calcite. *Geol.Soc.Amer.Bull.*, 101: 377-390.

Brass G.W. (1976) The variation of the marine  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio during Phanerozoic time: interpretation using a flux model. *Geochim. Cosmochim. Acta* 40, 721-730.

Chaisi, S.D. and Schmitz, B. (1995) Stable ( $\delta^{13}\text{C}$ ,  $\delta^{18}\text{O}$ ) and strontium ( $^{87}\text{Sr}/^{86}\text{Sr}$ ) isotopes through the Paleocene at Gebel Aweina, eastern Thetyan region. *Paleogeogr. Paleoclimatol. Paleocol.* 116, 103-129.

Cummins, D.I. and Elderfield, H. (1994) The strontium isotopic composition of Brigantian /late Dinantian seawater. *Chem. Geol.* 118, 255-270.

Dash, E.J. and Biscaye, P.E. (1971) Isotope composition of strontium in Cretaceous-Recent pelagic foraminifera. *Earth Planet. Sci. Lett.* 11, 201-204.

Denison, R.E.; Koepnick, R.B.; Burke, W.H.; Hetherington, E.A.; Fletcher, A. (1994): Construction of the Mississippian, Pennsylvanian and Permian seawater  $^{87}\text{Sr}/^{86}\text{Sr}$  curve. - *Chemical Geology (Isotope Geoscience Section)* 112, 145-167.

DePaolo, D.J. and Ingram, B.L. (1985) High-Resolution stratigraphy with Strontium isotopes. *Science* 227, 938-941.

Dia, A.N., Cohen, A.S., O'Nions, R.K. and Shackelton, N.J. (1992) Seawater Sr isotope variation over the past 300 kyr and influence of global climate cycles. *Nature* 356, 786-788.

Farrell, J. W., Clemens, S. C. and Gromet, L. P. (1995). Improved chronostratigraphic reference curve of late Neogene seawater  $^{87}\text{Sr}/^{86}\text{Sr}$ . *Geology*, 23: 403-406.

Hess, J. L., Scott, D., Bender, M. L., Kennett, J. P. and Schilling, J. -G., 1989. The Oligocene marine microfossil record: Age assessments using strontium isotopes. *Paleoceanography*, 4, 6, 655-679.

Hess, J., Bender, M.L. and Schilling, J.-G. (1986) Evolution of the ratio of Strontium-87 to Strontium-86 in seawater from Cretaceous to present. *Science* 231, 979-984.

Hodell, D.A. and Woodruff, F. (1994) Variations in strontium isotopic ratio of the seawater during the Miocene: Stratigraphic and geochemical implications. *Paleoceanography* 9, 405-426.

Hodell, D.A., Mead, G.A., and Mueller, P.A. (1990) Variation in the strontium isotopic composition of seawater (8 Ma to present): Implications for chemical weathering rates and dissolved fluxes to the oceans. *Chem. Geol.*, 80, 291-307.

Hodell, D.A., Mueller, P.A. and Garrido, J.R. (1991) Variations in the strontium isotopic composition of seawater during the Neogene. *Geology* 19, 24-27.

Hodell, D.A., Mueller, P.A., McKenzie, J.A. and Mead, G.A. (1989) Strontium isotope stratigraphy and geochemistry of the late neogene ocean. *Earth Planet. Sci. Lett.* 92, 165-178.

Jones, C.J., Jenkyns, H.C., Coe, A.L. and Hesselbo, S. (1994a) Strontium isotopic variations in the Jurassic Cretaceous seawater. *Geochim. Cosmochim. Acta.* 58, 3061-3074.

Jones, C.E., Jenkyns, H.C., Hesselbo, S.P. (1994b) Strontium isotopes in the Early Jurassic seawater. *Geochim. Cosmochim. Acta.* 58, 1285-1305.

Koepnick R.B., Burke W.H., Denison R.E., Heatherington E.A., Nelson H.F., Otto J.B. and Waite L.E. (1985) Construction of the Seawater  $^{87}\text{Sr}/^{86}\text{Sr}$  curve for the Cenozoic and Cretaceous: supporting data. *Chem. Geol. Isot. Geosc. Sect.* 58, 55-81.

Kürschner, W., Becker, R.T., Buhl, D and Veizer, J. 1993. Strontium isotopes in conodonts: Devonian-Carboniferous transition, the northern Rhenish Slate Mountains, Germany. *Ann.Géol.Belg.*, 155: 595-622

Martin, E.E. and McDougall, J.D. (1991) Seawater Sr isotopes at the Cretaceous/Tertiary boundary. *Earth Planet. Sci. Lett.* 104, 166-180.

Martin, E.E.; Macdougall, J.D. (1995): Sr and Nd isotopes at the Permian/Triassic boundary: A record of climate change. *Chemical Geology*, 125: 73-99.

McArthur, J.M., Kennedy, W.J., Chen, M., Thirlwall, M.F. and Gale, A.S. (1994) Strontium isotope stratigraphy for Late Cretaceous time: Direct numerical calibration of the Sr isotope curve based on the US Western Interior. *Palaeogr., Palclimat., Paleoecol.*, 108, 95-119.

McKenzie J.A., Isern A., Elderfield H, Williams A and Swart P.K. (1993) 33. Strontium Isotope dating of paleoceanographic, lithologic, and dolomitization events on the northeastern Australian margin, Leg 133. *Proc. of the ODP, Sci. Res.* 133, 489-498.

McKenzie, J.A., Hodell, D.A., Mueller, P.A. and Mueller, D.W. (1988) Application of strontium isotopes to late Miocene-early Pliocene stratigraphy. *Geology* 16, 1022-1025.

Oslick, J.S., Miller, K.G., Feigenson, M.D. and Wright, J.D. (1994) Oligocene-Miocene strontium isotopes: Stratigraphic revisions and correlations to an inferred glacioeustatic record. *Paleoceanography*, 9, 327-443.

Peterman, Z.E., Hedge, C.E. and Tourtelot, H.A. (1970) Isotopic composition of strontium in seawater throughout Phanerozoic time. *Geochim. Cosmochim. Acta* 34, 105-120.

Popp, N.B., Podosek, F.A., Brannon, J.C., Anderson, T.F. and Pier, J. 1986c.  $^{87}\text{Sr}/^{86}\text{Sr}$  in Permo-Carboniferous seawater from the analyses of well preserved brachiopod shells. *Geochim.Cosmochim.Acta*, 50: 1321-1328.

Quing, H, Barnes C. R., Buhl, D., and Veizer, J., The Sr isotopic composition of Ordovician and Silurian brachiopods and conodonts: relationships to geological events and implications for coeval seawater. *Geochim. et Cosmochim. Acta* (submitted).

Quinn T.M., Taylor F.W. and Halliday A.N. (1994) 6. Strontium-Isotopic dating of neritic carbonates and Bougainville Guyot (Site 831), New Hebrides island arc. *Proc. of ODP, Sci. Res.* 134, 89-95.

Ruppel, S. C., James, E. W., Barrick, J. E., Nowlan, G., and Uyeno, T. T., 1996, High-resolution  $^{87}\text{Sr}/^{86}\text{Sr}$  chemostratigraphy of the Silurian: Implications for event correlation and strontium flux: *Geology*, v. 24, p. 831-834.

Veizer, J.; Compston, W. (1974):  $^{87}\text{Sr}/^{86}\text{Sr}$  composition of seawater during the Phanerozoic. - *Geochimica et Cosmochimica Acta* 38, 1461-1484.

Wadleigh M.A., Veizer J. (1992)  $^{18}\text{O}/^{16}\text{O}$  and  $^{13}\text{C}/^{12}\text{C}$  in Lower Paleozoic brachiopods: isotopic composition of sea water. *Geochim. et Cosmochim. Acta* 56, 431-443.

Wenzel, B., 1997, Isotopenstratigraphie silurischer Abfolgen und deren palaeozeanographische Interpretation: *Erlanger Geologische Abhandlungen*, v. 129, 117 p.

Whittaker S.G. and Kyser T.K. (1993) Variations in the neodymium and strontium isotopic composition and REE content of molluscan shells from the Cretaceous Western Interior seaway. *Geochim et Cosmochim. Acta* 57, 4003-4014.