

**Brief overview on Grothendieck-Serre's conjecture** The following conjecture was stated by A. Grothendieck in 60's:

**Conjecture.** *Let  $G$  be a smooth reductive group scheme over a local regular ring  $R$  and  $K$  be the field of fractions of  $R$ . Then the induced map*

$$H_{et}^1(R, G) \rightarrow H_{et}^1(K, G)$$

*has trivial kernel.*

Observe that the pointed set  $H_{et}^1(R, G)$  classifies principal  $G$ -bundles over  $R$ , i.e., the conjecture says that any principal  $G$ -bundle which is rationally trivial is (Zariski-)locally trivial.

If  $R$  is of geometric type over a field  $k$  and  $G$  is defined over  $k$ , then the conjecture is known as Serre's Conjecture. If  $k$  is infinite, the latter was proven by Colliot-Thelene, Ojanguren [CO92] and Raghunathan [Ra94]. The case of a finite field  $k$  remains open till now.

The Grothendieck's conjecture was proven in the following cases:

- E. Nisnevich
  - $\dim R = 1$  or –  $R$  is Henselian of arbitrary dimension [Ni84]
  - $\dim R = 2$  and  $G$  is quasi-split over  $R$  [Ni89]
- J.-L. Colliot-Thelene, J.-J. Sansuc [CS87]
  - $G$  is an arbitrary torus over  $R$ .
- M. Ojanguren
  - $R$  is of geometric type and  $G = O_q$  is the orthogonal group of a non-degenerate quadratic form  $q$  over  $R$  [Oj80].
  - $\dim R = 2$  and  $G = O_q$  [Oj82].
  - $\dim R = 2$  and  $G = SL_A$  is the special linear group associated to an Azumaya algebra  $A$  over  $R$  [Oj82].
- I. Panin, A. Suslin [PS97]
  - $R$  is of geometric type and  $G = SL_A$
- M. Ojanguren, I. Panin [OP01]
  - $R$  contains a field of  $char \neq 2$  and  $G = U_{A,\sigma}$  is the unitary group of an Azumaya algebra  $A$  over  $R$  with involution  $\sigma$  of the second kind.
  - $R$  contains a field of  $char \neq 2$  and  $G = O_{A,\sigma}$  is the orthogonal group of an Azumaya algebra with involution  $\sigma$  of the first kind.
- K. Zainoulline [Za01]

- $R$  contains an infinite field of  $\text{char} \neq 2$  and  $G = SU_{A,\sigma}$  is the special unitary group associated to an Azumaya algebra  $A$  over  $R$  with an involution  $\sigma$  of the second kind.
  - The case  $R = SL_A$  for any  $R$  containing a field.
  - M. Ojanguren, I. Panin, K. Zainoulline [OPZ04]
    - $R$  contains an infinite field of  $\text{char} \neq 2$  and  $G = Spin_q$  is the Spinor group associated to a non-degenerate quadratic form over  $R$ .
  - I. Panin [Pa05]
    - $G$  is any adjoint group of classical type.
  - V. Chernousov, I. Panin [CP07]
    - $R$  contains a field of  $\text{char} \neq 2$  and  $G$  is of type  $G_2$  over  $R$ .
- To finish the case of groups of classical types it remains to prove
- $G = Spin_{A,\sigma}$ , where  $A$  is an Azumaya algebra over  $R$  with orthogonal involution  $\sigma$ ,  $\text{ind}(A) \geq 2$ .

## References

- [CP07] Chernousov, V., Panin, I. Purity for  $G_2$ -torsors. *Comptes Rendus Mathematique* 345 (2007), no. 6, 307–312.
- [CO92] Colliot-Thelene, J.-L.; Ojanguren, M. Espaces Principaux Homogenes Localement Triviaux. *Publ. Math. IHES* 75 (1992), 97–122.
- [CS87] Colliot-Thelene, J.-L.; Sansuc, J.-J. Principal Homogeneous Spaces under Flasque Tori: Applications. *J. of Algebra* 106 (1987), 148–205.
- [Ni89] Nisnevich, Y. Rationally trivial principal homogeneous spaces, purity and arithmetic of reductive group schemes over extensions of two-dimensional local regular rings. *C. R. Acad. Sci. Paris, t.309, Serie I* (1989), 651–655.
- [Ni84] Nisnevich, Y. Rationally Trivial Principal Homogeneous Spaces and Arithmetic of Reductive Group Schemes Over Dedekind Rings. *C. R. Acad. Sc. Paris, t. 299, Serie I, n.1* (1984), 5–8.
- [OPZ04] Ojanguren, M.; Panin, I.; Zainoulline, K. On the norm principle for quadratic forms. *J.Ramanujan Math. Soc.* 19 (2004), No.4, 1-12.
- [OP01] Ojanguren, M. and Panin, I. Rationally trivial Hermitian spaces are locally trivial. *Math. Z.* 237 (2001), pp. 181–198.

- [Oj82] Ojanguren, M. Unités représentées par des formes quadratiques ou par des normes réduites, in Algebraic K-theory, Oberwolfach, 1980, t. II, L.N.M., 967, 291-299, Springer-Verlag, 1982.
- [Oj80] Ojanguren, M. Quadratic forms over regular rings, Journal of the Indian Math. Soc., 44 (1980), 109–116.
- [Pa05] Panin, I. Purity for multipliers. Algebra and number theory, 66–89, Hindustan Book Agency, Delhi, 2005.
- [PS97] Panin, I. A.; Suslin, A. A. On a conjecture of Grothendieck concerning Azumaya algebras. (Russian) Algebra i Analiz 9 (1997), no. 4, 215–223, translation in St. Petersburg Math. J. 9 (1998), no. 4, 851–858.
- [Ra94] Raghunathan, M.S. Principal bundles admitting a rational section, Invent. math. 116 (1994), 409–423.
- [Za01] Zainoulline, K. On Grothendieck’s Conjecture about principal homogeneous spaces for some classical algebraic groups. St.Petersburg Math. J. 12 (2001), no.1, 117-143.