

Negative Frequency-Dependent Selection of Sexually Antagonistic Alleles in *Myodes glareolus* Mikael Mokkonen, *et al. Science* **334**, 972 (2011); DOI: 10.1126/science.1208708

This copy is for your personal, non-commercial use only.

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by clicking here.

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines here.

The following resources related to this article are available online at www.sciencemag.org (this infomation is current as of November 18, 2011):

Updated information and services, including high-resolution figures, can be found in the online version of this article at: http://www.sciencemag.org/content/334/6058/972.full.html

Supporting Online Material can be found at: http://www.sciencemag.org/content/suppl/2011/11/16/334.6058.972.DC1.html

This article **cites 27 articles**, 6 of which can be accessed free: http://www.sciencemag.org/content/334/6058/972.full.html#ref-list-1

This article appears in the following **subject collections:** Evolution http://www.sciencemag.org/cgi/collection/evolution

Negative Frequency-Dependent Selection of Sexually Antagonistic Alleles in *Myodes glareolus*

Mikael Mokkonen,¹* Hanna Kokko,² Esa Koskela,³ Jussi Lehtonen,^{2,4} Tapio Mappes,¹ Henna Martiskainen,³ Suzanne C. Mills⁵

Sexually antagonistic genetic variation, where optimal values of traits are sex-dependent, is known to slow the loss of genetic variance associated with directional selection on fitness-related traits. However, sexual antagonism alone is not sufficient to maintain variation indefinitely. Selection of rare forms within the sexes can help to conserve genotypic diversity. We combined theoretical models and a field experiment with *Myodes glareolus* to show that negative frequency-dependent selection on male dominance maintains variation in sexually antagonistic alleles. In our experiment, high-dominance male bank voles were found to have low-fecundity sisters, and vice versa. These results show that investigations of sexually antagonistic traits should take into account the effects of social interactions on the interplay between ecology and evolution, and that investigations of genetic variation should not be conducted solely under laboratory conditions.

The direction of evolutionary change generally cannot be predicted without taking into account interactions between conspecifics, be they competitors or mates (1, 2). Additionally, the fitness effects of particular genotypes can often be highly sex-specific (3). Alleles beneficial to a son's reproduction can be detrimental for a daughter's success, or vice versa, resulting in different optimal trait values (optima) between the sexes (4-8). The presence of these sexually antagonistic (SA) alleles in a variety of organisms is now widely acknowledged (9-11), but less is known about how variation in such alleles is maintained. All else being equal, SA selection does not decrease genetic variation as rapidly as selection that favors the same alleles in both males and females (6, 12).

However, SA selection is not as beneficial for genetic diversity as true negative frequencydependent maintenance of alternative alleles, whereby rare genotypes are favored in reproduction (13). The process of variation depletion may end in intralocus sexual conflict that remains unresolved, such that the traits of males, females, or both do not match their optima [leading to a cost of SA alleles, the gender load (14)]. It may also end in resolved conflict if gene expression evolves to become more sex-specific (15). In the former case sexual antagonism is preserved, in the latter case it disappears, but variation as such is lost in both. Because gender load can exist even if all individuals possess the same genotype—that is, the one that is the best compromise between male and female fitness variation is not necessarily maintained by gender load per se. Sexual antagonism, however, becomes potentially powerful for variance maintenance once it combines with sex linkage (16), maternal effects (17), or assortative mating (18). Here, we focused on the role of frequency-dependent selection as an alternative for making sexual antagonism successful at maintaining genetic variation.

We tested whether frequency-dependent selection on dominance in males and/or females can maintain SA variation in field populations of a common European mammal, the bank vole (*Myodes glareolus*). The reproductive effort of bank voles is negatively frequency-dependent in the field (19), attesting to the direct influence of an individual's neighbors on the population dynamics and life history evolution in this species. Previous experiments with this species have also shown that testosterone is under sexual and SA selection and is the most important determinant of dominance in male-male competition (20-22). This hormone also affects a variety of evolutionarily important processes such as spermatogenesis (23), immune function (24, 25), and secondary sex trait growth (26).

A haploid model with intralocus sexual conflict has shown that genetic variation can be maintained if the antagonism is sufficiently "balanced"-that is, if the relative fitness differences within males and females are of similar magnitude (27). This can lead to cyclic dynamics of morph frequency (28) or a protected polymorphism (29) where allele frequencies are roughly invariant over time. The maintenance of polymorphisms can be facilitated if the SA alleles are sex-linked (16), or if the fitness costs of SA alleles are nearly neutral when averaged between sexes and over their lifetimes (30). If we assume additional negative effects for each competitive type as their frequency becomes stronger, maintaining SA genetic variation becomes considerably easier.

Our integrated theoretical-empirical study included a large-scale field experiment, replicated over 2 years with a total of 31 populations, that exposed bank voles to terrestrial and avian predators as well as naturally occurring weather conditions and food resources (31). We first artificially selected groups in the laboratory (lines) to create behaviorally dominant males with sisters of low fecundity, and vice versa (31). In a series of malemale competition trials, two males of opposing behavioral dominance competed with each other to mate with a wild female in estrus. The highdominance males overwhelmingly outcompeted the low-dominance males in mating success trials $(\chi^2 = 59.71, N = 168, P < 0.001)$ (Fig. 1A), and they had a significantly higher plasma testosterone level (generalized linear mixed model; line: $F_{1,161} =$ 6.18, P = 0.014) (Fig. 1B). We then assigned males and females to large outdoor field enclosures by manipulating the reproductive quality (high or low behavioral dominance for males, high or



Fig. 1. Experimental data for males artificially selected for high and low dominance. (**A**) Relative mating success in laboratory male-male competition for a female in estrus. (**B**) Mean plasma testosterone level. (**C**) Mean frequency dependence of reproductive success in the field. Post hoc testing showed that the number of offspring sired by rare dominant males was significantly greater than the number sired by common dominant males (z = 2.97, P = 0.015). No other pairwise groupings differed significantly from each other (P > 0.1). Mf (black bars), high-dominance males with low-fecundity sisters; mF (white bars), low-dominance males with high-fecundity sisters; R, rare; C, common. Error bars in (B) and (C) denote SEM.

¹Centre of Excellence in Evolutionary Research, Department of Biological and Environmental Science, University of Jyväskylä, 40014 Jyväskylä, Finland. ²Centre of Excellence in Biological Interactions, Division of Evolution, Ecology and Genetics, Research School of Biology, Australian National University, Canberra, ACT 0200, Australia. ³Department of Biological and Environmental Science, University of Jyväskylä, 40014 Jyväskylä, Finland. ⁴Department of Biological and Environmental Sciences, University of Helsinki, 00014 Helsinki, Finland. ⁵Laboratoire d'Excellence "CORAIL," USR 3278 CRIOBE, CNRS-EPHE, CBETM de l'Université de Perpignan, 66860 Perpignan Cedex, France.

^{*}To whom correspondence should be addressed. E-mail: mikael.mokkonen@gmail.com



Fig. 2. The outcome of the model, expressed as the frequency of the *A* allele after convergence for three different frequency dependence scenarios. Allele frequencies smaller than 1 indicate that variation is maintained. (**A**) No explicit frequency dependence. (**B**) Explicit negative frequency dependence of SA alleles in all male genotypes. (**C**) Explicit negative frequency dependence only in the behaviorally dominant *AA* males. *AA*, *Aa*, and *aa* males have relative mating success values of $\alpha 1$, $\alpha 2$, and $\alpha 3$, and maximum frequency dependence values of $\beta 1$, $\beta 2$, and $\beta 3$. *D*_{male} and *D*_{female} represent

genetic dominance in males and females; γ and τ determine the shape of the frequency dependence curve [see (*31*) for details on the model and parameter estimates]. Parameter values used: $\alpha_1 = 80$, $\alpha_3 = 20$, $\beta_1 = 4.53$, $\beta_3 = 5.32$, $\gamma = 5$, $\tau = 0.5$, and (A) $\phi_1 = \phi_2 = \phi_3 = 0$, (B) $\phi_1 = 0.569$, $\phi_3 = 0.262$, (C) $\phi_1 = 0.569$, $\phi_2 = \phi_3 = 0$. The appropriate values for α_2 and β_2 [and ϕ_2 for the scenario in (B)], corresponding to different dominance values D_{male} and D_{female} , were calculated as $\alpha_2 = [D_{male}(\alpha_1 - \alpha_3) + \alpha_1 + \alpha_3]/2$, $\beta_2 = [D_{female}(\beta_3 - \beta_1) + \beta_3 + \beta_1]/2$, and $\phi_2 = [D_{male}(\phi_1 - \phi_3) + \phi_1 + \phi_3]/2$ according to Falconer's notation (*33*).

low fecundity for females) and frequency (rare or common) of male and female voles in these populations (fig. S1).

The reproductive success of males became negatively frequency-dependent in the field: Dominance was costly for males when it was the common tactic in the population (zero-inflated negative binomial count submodel; frequency, Z = -2.61, P = 0.009; line, Z = -1.01, P = 0.311; frequency × line, Z = 2.00, P = 0.046) (Fig. 1C). In contrast, the fitness of sisters demonstrated a sexually antagonistic effect without evidence for frequency dependence. The high-fecundity sisters of lowdominance males had significantly larger litter sizes than the sisters of high-dominance males (i.e., females selected for low fecundity) [5.32 \pm 0.24 versus 4.53 \pm 0.27 (SEM)], although, in a clear deviation from the male pattern, the reproductive output of these females did not depend on their frequency in the population (generalized linear model quasi-Poisson; frequency, t = 0.87, P = 0.389; line, t = 2.06, P = 0.043).

Combined with previous research on the selection of testosterone (20, 22, 25), our results suggest that there is potential for the sexes to experience antagonistic selection constraints in adaptation as a result of the physiological and behavioral components of their respective life history strategies. As shown in Fig. 1C, the male mating advantage is negatively frequency-dependent, which helps to explain why fixation of any allele is not predicted to occur.

In the next phase of our study, we extracted parameter estimates from our field data for use in our model (31), which expands previous work (27) to include diploid genetics and frequency dependence effects within males. Mating propensities of each male type either are constant or change with frequency; the latter choice creates implicit frequency dependence for the mating success of the other male type even if the mating propensity of this type was not set to depend on frequency, because an offspring sired by one male cannot be sired by another (females being a limiting resource). This can lead to the maintenance of genetic variability even if explicit frequency dependence is present in only one male type.

We ran our model with explicit frequency dependence on one or both male types (in the former case, frequency dependence is implicit for subordinate males), as well as with no frequency dependence, using parameters extracted from the experimental results. As we do not know the exact genetic system controlling the antagonistic traits in the study system, we ran the model using the entire range of genetic dominance parameters from full recessiveness to full dominance. In the absence of frequency dependence, genetic variation was maintained only when genetic dominance for both female and male traits was high (in other words, A is the dominant allele for males; a is dominant for females) (Fig. 2A). Frequency dependence of the magnitude we found for voles makes the maintenance of genetic variation much easier; it is maintained for a large proportion of the parameter range describing genetic dominance (Fig. 2B). It is maintained to a very similar degree even if frequency dependence arises only implicitly for low-dominance males (Fig. 2C). Our proposed mechanism thus appears reasonably robust, and future work could fruitfully combine these findings with other mechanisms known to promote maintenance of variation under SA selection (16-18).

Our results also suggest that studies of this type should not be restricted to laboratory environments. Seminatural conditions led to a sharp difference in the success of rare versus common behaviorally dominant males (Fig. 1C)—an effect that would have remained undetectable had our study been confined to the mating trials conducted in the laboratory. The difference in optimal food and housing conditions in the lab and the limitations of the field environment can have implications for physiological and behavioral measures (32) and fitness consequences alike. Thus, our theoretical and empirical results together show that selection favoring rare male morphs can maintain genetic variation in sexually antagonistic traits, while also indicating that ecological and social environments are important in defining the trait optima for males and females.

References and Notes

- 1. P. Bijma, J. Evol. Biol. 23, 194 (2010).
- 2. H. Kokko, A. López-Sepulcre, Ecol. Lett. 10, 773 (2007).
- 3. A. Y. K. Albert, S. P. Otto, *Science* **310**, 119 (2005). 4. G. A. Parker, in *Sexual Selection and Reproductive*
- Competition in Insects, M. S. Blum, N. A. Blum, Eds. (Academic Press, London, 1979), pp. 123–166.
- 5. W. R. Rice, Science 256, 1436 (1992).
- 6. K. Foerster et al., Nature 447, 1107 (2007).
- 7. K. M. Fedorka, T. A. Mousseau, Nature 429, 65 (2004).
- M. Delcourt, M. W. Blows, H. D. Rundle, Proc. Biol. Sci. 276, 2009 (2009).
- T. Connallon, R. M. Cox, R. Calsbeek, *Evolution* 64, 1671 (2010).
- 10. R. M. Cox, R. Calsbeek, Am. Nat. 173, 176 (2009).
- 11. G. Arnqvist, L. Rowe, *Sexual Conflict* (Princeton Univ. Press, Princeton, NJ, 2005).
- 12. M. D. Hall, S. P. Lailvaux, M. W. Blows, R. C. Brooks, *Evolution* 64, 1697 (2010).
- F. J. Ayala, C. A. Campbell, Annu. Rev. Ecol. Syst. 5, 115 (1974).
- S. Bedhomme, A. K. Chippindale, in *Sex, Size and Gender Roles: Evolutionary Studies of Sexual Size Dimorphism*,
 D. J. Fairbairn, W. U. Blankenhorn, T. Székely, Eds. (Oxford Univ. Press, New York, 2007), pp. 185–194.
- 15. G. S. van Doorn, Ann. N.Y. Acad. Sci. 1168, 52 (2009).
- 16. W. R. Rice, Evolution 38, 735 (1984).
- 17. M. M. Patten, D. Haig, Biol. Lett. 5, 667 (2009).
- 18. G. Arngvist, Evolution 65, 2111 (2011).
- 19. T. Mappes et al., PLoS One 3, e1687 (2008).
- 20. S. C. Mills et al., Am. Nat. 173, 475 (2009).
- 21. S. C. Mills, A. Grapputo, E. Koskela, T. Mappes, *Proc. Biol. Sci.* **274**, 143 (2007).
- M. Mokkonen, E. Koskela, T. Mappes, S. C. Mills, J. Anim. Ecol. 10.1111/j.1365-2656.2011.01903.x (2011).
- T. R. Birkhead, D. J. Hosken, S. Pitnick, Eds., Sperm Biology: An Evolutionary Perspective (Academic Press, Oxford, 2009).
- 24. I. Folstad, A. J. Karter, Am. Nat. 139, 603 (1992).
- 25. E. Schroderus et al., Am. Nat. 176, E90 (2010).
- M. Zuk, T. Johnsen, T. Maclarty, Proc. Biol. Sci. 260, 205 (1995).
- 27. H. Kokko, R. Brooks, Ann. Zool. Fenn. 40, 207 (2003).
- 28. B. Sinervo, R. Calsbeek, Annu. Rev. Ecol. Syst. 37, 581 (2006).
- E. I. Svensson, J. Abbott, R. Härdling, Am. Nat. 165, 567 (2005).

- A. K. Chippindale, J. R. Gibson, W. R. Rice, Proc. Natl. Acad. Sci. U.S.A. 98, 1671 (2001).
- 31. See supporting material on Science Online.
- R. M. Calisi, G. E. Bentley, *Horm. Behav.* 56, 1 (2009).
- D. S. Falconer, T. F. C. MacKay, *Introduction to Quantitative Genetics* (Pearson Prentice Hall, London, ed. 4, 1996).
- Acknowledgments: Supported by Academy of Finland grants 115961, 119200, and 218107 (E.K.), 132190 (T.M.), and 103508 and 108566 (S.C.M.); the Vanamo Biological Society and Ehrnrooth Foundation (M.M.); the Australian Research Council and Australian National

University (H.K. and J.L.); the Finnish Cultural Foundation and Emil Aaltonen Foundation (J.L.); and the Centre of Excellence in Evolutionary Research, University of Jyväskylä. We thank the staff of the Experimental Animal Unit and Konnevesi Research Station, University of Jyväskylä; R. Närä and H. Pietiläinen for logistical support; T. Laaksonen, V. Lummaa, and three anonymous reviewers for comments; and C. Soulsbury for statistical advice. The authors declare no conflicts of interest. All co-authors designed this study; M.M., E.K., T.M. and H.M. collected and analyzed the empirical data; J.L. analyzed theoretical results with input from all authors; and M.M. led the preparation of the manuscript with input from all co-authors. Authors after the first author are listed in alphabetical order. Data have been deposited in the Dryad Repository (doi:10.5061/dryad.6m0f6870).

Supporting Online Material

www.sciencemag.org/cgi/content/full/334/6058/972/DC1 Materials and Methods Figs. S1 to S3 Table S1 References (*34–41*)

cluster is referred to as the P cluster. The P clus-

20 May 2011; accepted 29 September 2011 10.1126/science.1208708

X-ray Emission Spectroscopy Evidences a Central Carbon in the Nitrogenase Iron-Molybdenum Cofactor

Kyle M. Lancaster,¹ Michael Roemelt,² Patrick Ettenhuber,² Yilin Hu,³ Markus W. Ribbe,³* Frank Neese,^{2,4}* Uwe Bergmann,⁵* Serena DeBeer^{1,4}*

Nitrogenase is a complex enzyme that catalyzes the reduction of dinitrogen to ammonia. Despite insight from structural and biochemical studies, its structure and mechanism await full characterization. An iron-molybdenum cofactor (FeMoco) is thought to be the site of dinitrogen reduction, but the identity of a central atom in this cofactor remains unknown. Fe K β x-ray emission spectroscopy (XES) of intact nitrogenase MoFe protein, isolated FeMoco, and the FeMoco-deficient Δ *nifB* protein indicates that among the candidate atoms oxygen, nitrogen, and carbon, it is carbon that best fits the XES data. The experimental XES is supported by computational efforts, which show that oxidation and spin states do not affect the assignment of the central atom to C⁴⁻. Identification of the central atom will drive further studies on its role in catalysis.

In trogenase (N₂ase), found in symbiotic and free-living diazotrophs, catalyzes the reduction of dinitrogen (N₂) to ammonia (NH₃) using eight electrons, eight protons, and 16 MgATPs (ATP, adenosine triphosphate) (1). Industrially, the same reaction is performed by the Haber-Bosch process that produces more than 100 million tons of NH₃ each year, thereby accounting for ~1.4% of global energy consumption. Understanding how nature activates the strongest homodinuclear bond in chemistry, the triple bond of N₂, is the key for the future design of molecular catalysts.

The high-resolution crystal structure of N_2 ase determined by Einsle *et al.* (2) showed that the active site of the molybdenum-iron (MoFe) protein component of N_2 ase binds a complex cluster consisting of seven iron ions, one molybde-

*To whom correspondence should be addressed: mribbe@uci. edu (M.W.R.); frank.neese@mpi-mail.mpg.de (F.N.); bergmann@ slac.stanford.edu (U.B.); serena.debeer@mpi-mail.mpg.de (S.D.) num ion, and nine sulfides (Fig. 1A); this cluster is referred to as the iron-molybdenum cofactor (FeMoco) and is thought to be the site of dinitrogen activation. For each FeMoco (of which there are two in the $\alpha_2\beta_2$ tetrameric MoFe protein) there is an additional cluster that consists of eight irons and seven sulfides (Fig. 1B); this

ters serve as electron-transfer sites. Several reaction intermediates in nitrogenase catalysis have recently been observed (3, 4). However, despite the progress in the experimental and theoretical analysis of the FeMoco (4-7), neither the reaction that occurs at the FeMoco nor the structure of FeMoco has been fully clarified. In 2002, Einsle et al. identified a light atom in the center of FeMoco that could be attributed to a single, fully ionized C, N, or O atom (2). No consensus has since emerged concerning the nature of this key atom. Study of FeMoco by electron paramagnetic resonance and related techniques is complicated by complex spin-couplings between the open-shell ions, which are not fully understood. Mössbauer spectroscopy suffers from spectral crowding, and neither nuclear resonance vibrational spectroscopy nor extended x-ray absorption fine structure are sufficiently conclusive (8).

Herein, we report iron K β valence-to-core (V2C) x-ray emission spectroscopy (XES) of N₂ase and demonstrate that these data provide a signature for the presence and identity of the central atom. K α and K β XES monitor the emission of photons after ionization of a metal 1s electron. The K $\beta_{1,3}$ emission line (~7040 to 7070 eV) corresponds to an electric dipole allowed $3p \rightarrow 1s$ transition. To higher emission energies,



Fig. 1. The FeMoco (**A**) and P cluster (**B**) of nitrogenase (adapted from the Protein Data Bank: identification number 1MIN). Orange, Fe; yellow, S; light blue, Mo; black, C^{4-} , N³⁻, or O²⁻; dark blue, nitrogen; gray, carbon. For clarity, the homocitrate and histidine ligands to the Mo have been omitted.

¹Department of Chemistry and Chemical Biology, Cornell University, Ithaca, NY 14853, USA. ²Institut für Physikalische und Theoretische Chemie, Universität Bonn, D-53115 Bonn, Germany. ³Department of Molecular Biology and Biochemistry, University of California, Irvine, CA 92697, USA. ⁴Max-Planck-Institut für Bioanorganische Chemie, Stiftstrasse 34-36, D-45470 Mülheim an der Ruhr, Germany. ⁵Linac Coherent Light Source, SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA.