

Invariant bilinear forms of algebras given by faithfully flat descent

E. Neher

*Department of Mathematics and Statistics
University of Ottawa
Ottawa, ON, Canada K1N 6N5
neher@uottawa.ca*

A. Pianzola

*Department of Mathematics, University of Alberta
Edmonton, AB, Canada T6G 2G1
Centro de Altos Estudios en Ciencias Exactas
Avenida de Mayo 866, (1084) Buenos Aires, Argentina
a.pianzola@gmail.com*

D. Prelat* and C. Sepp†

*Centro de Altos Estudios en Ciencias Exactas
Avenida de Mayo 866, (1084) Buenos Aires, Argentina
*dprelat@caece.edu.ar
†csepp@caece.edu.ar*

Received 8 August 2013

Accepted 7 November 2013

Published 30 December 2013

The existence of nondegenerate invariant bilinear forms is one of the most important tools in the study of Kac–Moody Lie algebras and extended affine Lie algebras. In practice, these forms are created, or shown to exist, either by assumption or in an *ad hoc* basis. The purpose of this work is to describe the nature of the space of invariant bilinear forms of certain algebras given by faithfully flat descent (which includes the affine Kac–Moody Lie algebras, as well as Azumaya algebras and multiloop algebras) within a functorial framework. This will allow us to conclude the existence, uniqueness and nature of invariant bilinear forms for many important classes of algebras.

Keywords: Invariant bilinear form; twisted form; centroid; faithfully flat descent.

Mathematics Subject Classification 2010: 17B67, 17B01, 12G05, 20G10

1. Introduction

One of the key ingredients for the study of Kac–Moody Lie algebras is the (generalized) Casimir operator. The existence of this operator is in turn based upon the existence of a symmetric invariant nondegenerate bilinear form on the Lie algebra.

Kac–Moody Lie algebras admitting such bilinear forms are called symmetrizable, and the most important example of these is given by the affine Kac–Moody Lie algebras.

It is also known [26] that the affine algebras are precisely the twisted forms (given by Galois, hence also faithfully flat descent) of the “split” affine algebras, namely of algebras of the form $\mathfrak{g} \otimes_k k[t^{\pm 1}]$ where \mathfrak{g} is a finite-dimensional simple Lie algebra over an algebraically closed field k of characteristic 0. In this paper we shall establish the representability (in a functorial sense) and explicit description of the space of invariant k -bilinear forms for a large class of (in general infinite dimensional) algebras given by faithfully flat descent.

The techniques developed in this paper not only apply to Lie algebras, but to other classes of algebras as well, such as Azumaya algebras, octonion, alternative or Jordan algebras; see Sec. 6 for an (incomplete) list of examples. Just as in the theory of Kac–Moody Lie algebras, the existence of a nondegenerate or even nonsingular invariant bilinear form on these algebras has important structural consequences. We therefore follow the approach of [23] and study invariant bilinear forms of arbitrary (nonassociative) algebras.

Among other things we will recover [18, Lemma 2.3], which considers this question for Lie algebras in the untwisted case. The need to consider graded invariant forms in the study of extended affine Lie algebras will require a “graded version” of our main results. This will be given towards the end of the paper in Sec. 7, where we will provide, among other things, a classification-free proof of Yoshii’s theorem [32] for multiloop Lie tori stating that graded invariant bilinear forms are unique up to scalars. We will also obtain new (as well as shed new light on known) results on Azumaya algebras (Theorem 6.10).

The technique that we use to describe the nature of invariant bilinear forms of algebras is based on two crucial ingredients:

- That invariant bilinear forms are functorial in nature and that this functor is representable.
- Descent theory.

This leads us to outline a general theory of descent within a functorial setting which we find to be of independent interest. It is developed in Sec. 2 and later applied to the functor **IBF** representing invariant bilinear forms.

2. Descent for Functors Stable Under Base Change

The goal of this section is to show that functors from the category of algebras to the category of modules that are stable under base change preserve descended forms. Relevant to our paper is the special case of the “functor of invariant bilinear forms” **IBF** as stated in Corollary 4.4. To write the proof in this particular case, however, obscures the more general nature of the construction that is taking place. We have thus chosen a framework that allows the essence of the argument to come across.

Our setting, while somewhat abstract, is sufficiently ample for our purposes and of independent interest. By appealing to fibered categories, an even more general set up is possible in which the concept of functors stable under base change can be defined. It is nevertheless a delicate question to identify, within this more general setting, which arrows play the (crucial) role of faithfully flat base change and their accompanying descent theory. We leave it to the interested reader to explore such more general scenarios.

Definition 2.1 (Categories $k\text{-alg}$, $k\text{-ALG}$, $k\text{-MOD}$). We fix once and for all a commutative associative unital ring k .

We denote by $k\text{-alg}$ the category of commutative associative unital k -algebras with unital k -linear algebra homomorphisms as morphisms. The symbol $R \in k\text{-alg}$ means that R is an object in $k\text{-alg}$. We also use the notation R/k to describe this situation. By definition, R comes accompanied with a “structure” ring homomorphism $\sigma_{R,k} : k \rightarrow R$ under which R is viewed as a k -module.

Of course since R is a commutative ring, we also have the category $R\text{-alg}$. An object $S \in R\text{-alg}$ will be viewed as an object of $k\text{-alg}$ via $\sigma_{S,k} = \sigma_{S,R} \circ \sigma_{R,k}$. The arrows of $R\text{-alg}$ are then arrows of $k\text{-alg}$ and we have a natural forgetful functor $R\text{-alg} \rightarrow k\text{-alg}$.

Let $\alpha : R \rightarrow S$ be a morphism in $k\text{-alg}$, M be an R -module and N be an S -module. We say that a map $f : M \rightarrow N$ is an α -semilinear module homomorphism if it is additive and satisfies $f(rm) = \alpha(r)f(m)$ for all $r \in R$ and $m \in M$. Such a map f is necessarily k -linear, namely if M and N are viewed as k -modules by means of the structure maps $\sigma_{R,k}$ and $\sigma_{S,k}$ then $f(cm) = cf(m)$ for all $c \in k$ and $m \in M$. If M and N have algebra structures and f preserves multiplication, then we say that f is an α -semilinear algebra homomorphism.

Given an α -semilinear module homomorphism $f : M \rightarrow N$ as above and a morphism $\beta : S \rightarrow T$ in $k\text{-alg}$, there exists a unique β -semilinear module homomorphism $f \otimes \beta : M \otimes_R S \rightarrow N \otimes_S T$ satisfying $(f \otimes \beta)(m \otimes s) = f(m) \otimes \beta(s)$. It is clear that if f is an algebra homomorphism, then so is $f \otimes \beta$.

The objects of the category $k\text{-ALG}$ are pairs (R, A) consisting of an $R \in k\text{-alg}$ and an R -algebra A . By this we mean an R -module A together with an R -bilinear map $A \times A \rightarrow A$, $(a_1, a_2) \mapsto a_1 a_2$. In particular, we do not require any further identities (even though Lie algebras are our main interest, all algebras are considered). A *morphism* in $k\text{-ALG}$, written as $(\alpha, f) : (R, A) \rightarrow (S, B)$, is a pair consisting of a morphism $\alpha : R \rightarrow S$ in $k\text{-alg}$ together with an α -semilinear algebra homomorphism $f : A \rightarrow B$.^a

The category $k\text{-MOD}$ is defined in complete analogy to $k\text{-ALG}$, with algebras replaced by modules and algebra homomorphisms replaced by module homomorphisms. Thus, an object in $k\text{-MOD}$ is a pair (R, M) consisting of some $R \in k\text{-alg}$ and an R -module M and a morphism in $k\text{-MOD}$ is a pair $(\alpha, f) : (R, M) \rightarrow (S, N)$

^aThe category $k\text{-ALG}$ appears in similar form in [17, 3.2].

consisting of a morphism $\alpha: R \rightarrow S$ in $k\text{-alg}$ and an α -semilinear module homomorphism $f: M \rightarrow N$.

Definition 2.2 (Base change). Both categories $k\text{-ALG}$ and $k\text{-MOD}$ admit base change by objects of $k\text{-alg}$. We explain this for $k\text{-ALG}$ and at the same time also set up our notation. Let $\alpha: R \rightarrow S$ be a morphism in $k\text{-alg}$ and let A be an R -algebra. Viewing S as an R -algebra via α , the tensor product $A \otimes_R S$ is an S -algebra whose product is given by $(a_1 \otimes s_1)(a_2 \otimes s_2) = (a_1 a_2) \otimes (s_1 s_2)$ for $a_i \in A$ and $s_i \in S$. Since we will in this section repeatedly use different algebra homomorphisms between R and S , it will be useful to temporarily employ the more precise notation $A \otimes_\alpha S$ instead of the traditional $A \otimes_R S$. We will also appeal to this notation elsewhere in the paper whenever it is necessary to emphasize the structure map α . We define

$$\alpha^A: A \rightarrow A \otimes_\alpha S, \quad a \mapsto a \otimes 1_S,$$

and note that $(\alpha, \alpha^A): (R, A) \rightarrow (S, A \otimes_\alpha S)$ is a morphism in $k\text{-ALG}$. Moreover, for any morphism $(\alpha, f): (R, A) \rightarrow (S, B)$ in $k\text{-ALG}$ there exists a unique S -linear algebra homomorphism $f^\alpha: A \otimes_\alpha S \rightarrow B$ satisfying $f^\alpha(a \otimes s) = sf(a)$. It is clear from the definitions that

$$\begin{array}{ccc} (R, A) & \xrightarrow{(\alpha, f)} & (S, B) \\ & \searrow (\alpha, \alpha^A) & \nearrow (\text{Id}_S, f^\alpha) \\ & & (S, A \otimes_\alpha S) \end{array} \tag{2.1}$$

is a commutative diagram in $k\text{-ALG}$. Assume that $\beta: S \rightarrow T$ is another morphism in $k\text{-alg}$. We can then add to the diagram (2.1) the morphism $(\beta, \beta^B): (S, B) \rightarrow (T, B \otimes_\beta T)$ and obtain the diagram

$$\begin{array}{ccc} (R, A) & \xrightarrow{(\alpha, f)} & (S, B) \\ (\alpha, \alpha^A) \downarrow & & \downarrow (\beta, \beta^B) \\ (S, A \otimes_\alpha S) & \xrightarrow{(\beta, f \otimes \beta)} & (T, B \otimes_\beta T) \end{array} \tag{2.2}$$

which commutes since $(\beta, f \otimes \beta) = (\beta, \beta^B) \circ (\text{Id}_S, f^\alpha)$. One should view this diagram as the base change of (α, f) by β .

Definition 2.3 (Functors over $k\text{-alg}$). The projection onto the first component defines functors $\Pi_A: k\text{-ALG} \rightarrow k\text{-alg}$ and $\Pi_M: k\text{-MOD} \rightarrow k\text{-alg}$. We say that a functor $F: k\text{-ALG} \rightarrow k\text{-MOD}$ is a functor over $k\text{-alg}$ if $\Pi_M \circ F = \Pi_A$, i.e.

$$\begin{array}{ccc} k\text{-ALG} & \xrightarrow{F} & k\text{-MOD} \\ & \searrow \Pi_A & \swarrow \Pi_M \\ & & k\text{-alg} \end{array}$$

is a commutative diagram. Such a functor F maps an object $(R, A) \in k\text{-ALG}$ to $F(R, A) = (R, F_R(A))$ for some R -module $F_R(A)$, and it sends a

morphism $(\alpha, f) : (R, A) \rightarrow (S, B)$ to $F(\alpha, f) = (\alpha, F_\alpha(f))$ for some α -semilinear map $F_\alpha(f) : F_R(A) \rightarrow F_S(B)$.

Convention. In what follows and without further explication, if $R = S$ and $\alpha = \text{Id}_S$ then we will simply write $F(f)$ instead of $F_\alpha(f)$.

Given a morphism $\alpha : R \rightarrow S$ in $k\text{-alg}$ and $(R, A) \in k\text{-ALG}$, we can apply F to the morphism $(\alpha, \alpha^A) : (R, A) \rightarrow (S, A \otimes_\alpha S)$ in $k\text{-ALG}$ of Definition 2.2 and get a morphism in $k\text{-MOD}$

$$F(\alpha, \alpha^A) = (\alpha, F_\alpha(\alpha^A)) : (R, F_R(A)) \rightarrow (S, F_S(A \otimes_\alpha S)).$$

Since this map is α -semilinear, it induces an S -linear map

$$\nu_{A,\alpha}^F : F_R(A) \otimes_\alpha S \rightarrow F_S(A \otimes_\alpha S), \quad m \otimes s \mapsto sF_\alpha(\alpha^A)(m) \tag{2.3}$$

of S -modules. These maps will play an essential role in our work for they constitute the essential ingredient in the definition of functors stable under base change. For convenience in what follows, if F is fixed in the discussion, we will denote $\nu_{A,\alpha}^F$ simply by $\nu_{A,\alpha}$.

Lemma 2.4. *Suppose that $F : k\text{-ALG} \rightarrow k\text{-MOD}$ is a functor over $k\text{-alg}$.*

(a) (F and ν commute) *Let $(\alpha, f) : (R, A) \rightarrow (S, B)$ be a morphism in $k\text{-ALG}$ and let $\beta : S \rightarrow T$ be a morphism in $k\text{-alg}$. Then the diagram*

$$\begin{array}{ccc} F_R(A) \otimes_\alpha S & \xrightarrow{F_\alpha(f) \otimes \beta} & F_S(B) \otimes_\beta T \\ \nu_{A,\alpha} \downarrow & & \downarrow \nu_{B,\beta} \\ F_S(A \otimes_\alpha S) & \xrightarrow{F_\beta(f \otimes \beta)} & F_T(B \otimes_\beta T) \end{array}$$

commutes.

(b) (Transitivity) *Let $R \xrightarrow{\alpha} S \xrightarrow{\beta} T$ be morphisms in $k\text{-alg}$ and let $(R, A) \in k\text{-ALG}$. Then the diagram*

$$\begin{array}{ccc} F_R(A) \otimes_\alpha S & \xrightarrow{\text{Id}_{F_R(A)} \otimes \beta} & F_R(A) \otimes_{\beta \circ \alpha} T \\ \nu_{A,\alpha} \downarrow & & \downarrow \nu_{A,\beta \circ \alpha} \\ F_S(A \otimes_\alpha S) & \xrightarrow{F_\beta(\text{Id}_A \otimes \beta)} & F_T(A \otimes_{\beta \circ \alpha} T) \end{array}$$

commutes.

Proof. (a) The maps $f \otimes \beta$ and $F_\alpha(f) \otimes \beta$ are described in Definitions 2.1 and 2.2. The bottom horizontal arrow is obtained by applying F to $(\beta, f \otimes \beta)$. For $m \in F_R(A)$ and $s \in S$ we have

$$\begin{aligned} (\nu_{B,\beta} \circ (F_\alpha(f) \otimes \beta))(m \otimes s) &= \beta(s)F_\beta(\beta^B)(F_\alpha(f)(m)), \text{ while} \\ (F_\beta(f \otimes \beta) \circ \nu_{A,\alpha})(m \otimes s) &= \beta(s)F_\beta(f \otimes \beta)(F_\alpha(\alpha^A)(m)). \end{aligned}$$

It therefore suffices to show that $F_\beta(\beta^B) \circ F_\alpha(f) = F_\beta(f \otimes \beta) \circ F_\alpha(\alpha^A)$. But this follows by applying the functor F to the commutative diagram (2.2).

(b) With the notation of (a) we have

$$(\nu_{A,\beta \circ \alpha} \circ (\text{Id} \otimes \beta))(m \otimes s) = \beta(s)F_{\beta \circ \alpha}((\beta \circ \alpha)^A)(m), \text{ while}$$

$$(F_\beta(\text{Id}_A \otimes \beta) \circ \nu_{A,\alpha})(m \otimes s) = \beta(s)F_\beta(\text{Id}_A \otimes \beta)F_\alpha(\alpha^A)(m).$$

It is therefore sufficient to show that $F((\beta \circ \alpha)^A) = F_\beta(\text{Id}_A \otimes \beta) \circ F_\alpha(\alpha^A)$. By functoriality, this is a consequence of $((\beta \circ \alpha), (\beta \circ \alpha)^A) = (\beta, \text{Id}_A \otimes \beta) \circ (\alpha, \alpha^A)$. \square

Definition 2.5 (Functors stable under base change). Let $F : k\text{-ALG} \rightarrow k\text{-MOD}$ be a functor over $k\text{-alg}$. We will say F is *stable under base change* if for all morphisms $\alpha \in k\text{-alg}$ and all $(R, A) \in k\text{-ALG}$ the S -module homomorphism $\nu_{A,\alpha}^F : F_R(A) \otimes_\alpha S \rightarrow F_S(A \otimes_\alpha S)$ defined in (2.3) is an isomorphism.

Example 2.6. An example of a functor stable under base change is the “invariant bilinear form functor” **IBF** of Definition 3.5, see Proposition 4.3(b), which is most relevant to our work. Of course the quintessential example of a functor $F : k\text{-ALG} \rightarrow k\text{-MOD}$ over $k\text{-alg}$ stable under base change is the “tensor product” functor that attaches to (R, A) the R -module $A \otimes_R A$ with the natural definition of F at the level of arrows. The stability of base change is given by the canonical S -module isomorphism $(A \otimes_R A) \otimes_R S \simeq (A \otimes_R S) \otimes_S (A \otimes_R S)$.

To justify the generality of this section we give another example. Let A be an R -algebra. Recall that its derived algebra is defined by

$$\mathcal{D}(A) = \text{Span}_R\{a_1 a_2 : a_i \in A\} = \text{Span}_\mathbb{Z}\{a_1 a_2 : a_i \in A\}.$$

If $(\alpha, f) : (R, A) \rightarrow (S, B)$ is a morphism in $k\text{-ALG}$, then $f(\mathcal{D}(A)) \subset \mathcal{D}(B)$. Hence by restriction we obtain a map $\mathcal{D}_\alpha(f) : \mathcal{D}(A) \rightarrow \mathcal{D}(B)$. If we view $\mathcal{D}(A)$ as a submodule of the R -module A it is immediate that $(\alpha, \mathcal{D}_\alpha(f)) : (R, \mathcal{D}(A)) \rightarrow (S, \mathcal{D}(B))$ is a morphism in $k\text{-MOD}$ and that the assignment

$$(R, A) \mapsto (R, \mathcal{D}(A)) \quad \text{and} \quad (\alpha, f) \mapsto (\alpha, \mathcal{D}_\alpha(f))$$

defines a functor $\mathcal{D} : k\text{-ALG} \rightarrow k\text{-MOD}$ over $k\text{-alg}$.

The explicit nature of the map $\nu_{A,\alpha}^\mathcal{D} : \mathcal{D}(A) \otimes_\alpha S \rightarrow \mathcal{D}(A \otimes_\alpha S)$ is clear: $a_1 a_2 \otimes s \mapsto (a_1 \otimes s)(a_2 \otimes 1) = a_1 a_2 \otimes s = (a_1 \otimes 1)(a_2 \otimes s)$. Thus $\nu_{A,\alpha}^\mathcal{D}$ is always surjective. This map, however, needs to be injective since there is no reason for the natural map $\mathcal{D}(A) \otimes_\alpha S \rightarrow A \otimes_\alpha S$ to be so (it is, for example, if $\alpha : R \rightarrow S$ is flat).

Consider a new functor $F : k\text{-ALG} \rightarrow k\text{-MOD}$ over $k\text{-alg}$ which assigns to (R, A) the pair $(R, A/\mathcal{D}(A))$ and is defined at the level of arrows in the natural way. We leave it to the reader to check that the surjectivity of $\nu_{A,\alpha}^\mathcal{D}$ easily implies that $\nu_{A,\alpha}^F$ is an isomorphism of S -modules. Thus F is stable under base change.

Remark 2.7. The results of this section can be generalized by replacing $k\text{-alg}$, $k\text{-ALG}$ and $k\text{-MOD}$ by subcategories stable under base change and by modifying the Definition 2.3 correspondingly. As we shall see, the most relevant case for us is that of functors which are stable under faithfully flat base change. We will leave it to the interested reader to work out the necessary axioms.

Descent Theory 2.8 (Faithfully flat descent of modules and algebras). We give a short review of the descent theory of modules and algebras. Our ultimate objective is to outline a descent theory in the setting of functors stable under faithfully flat base change. We also use the opportunity to introduce notation and a presentation of descent theory that is implicitly, but not explicitly used in the standard references ([13; 29; 30]), cf. [25]. Without these the formalism for descent in the functorial setting is impossible to redact.

Assume that S/R is faithfully flat. We let $S'' = S \otimes_R S$ and denote by $\alpha_i : S \rightarrow S''$, $i = 1, 2$, the “projections” defined by

$$\alpha_1(s) = s \otimes 1 \quad \text{and} \quad \alpha_2(s) = 1 \otimes s,$$

which allow us to view S'' as an S -algebra in two different ways. Note that since $\alpha : R \rightarrow S$ is faithfully flat, $\alpha_1 \circ \alpha = \alpha_2 \circ \alpha$. Suppose M and N are R -modules and that N is an S/R -form of M . Thus there exists an S -module isomorphism $\theta : (M \otimes_\alpha S) \rightarrow (N \otimes_\alpha S)$. To θ and $i = 1, 2$ we associate the S'' -module isomorphisms θ_i defined by the following commutative diagram:

$$\begin{array}{ccc} (M \otimes_\alpha S) \otimes_{\alpha_i} S'' & \xrightarrow[\simeq]{\theta \otimes_{\alpha_i} \text{Id}_{S''}} & (N \otimes_\alpha S) \otimes_{\alpha_i} S'' \\ \tau_i^M \downarrow \simeq & & \simeq \downarrow \tau_i^N \\ M \otimes_{\alpha_i \circ \alpha} S'' & \xrightarrow[\simeq]{\theta_i} & N \otimes_{\alpha_i \circ \alpha} S'' \end{array} \quad (2.4)$$

Here $\tau_i : S \otimes_{\alpha_i} S'' \rightarrow S''$ is defined by $\tau_i(s_1 \otimes s_2 \otimes s_3) = \alpha_i(s_1)(s_2 \otimes s_3)$, e.g., $\tau_2(s_1 \otimes s_2 \otimes s_3) = (s_2 \otimes s_1 s_3)$, while $\tau_i^M((m \otimes s_1) \otimes s_2 \otimes s_3) = m \otimes \alpha_i(s_1)(s_2 \otimes s_3)$. The maps τ_i^N are defined similarly. In what follows, τ_i^M and τ_i^N are viewed as S'' -linear maps.

The situation can be summarized by the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & M \simeq M \otimes_{\text{Id}_R} R & \xrightarrow{\text{Id}_M \otimes \alpha} & M \otimes_\alpha S & \begin{array}{c} \xrightarrow{\text{Id}_M \otimes \alpha_1} \\ \xrightarrow{\text{Id}_M \otimes \alpha_2} \end{array} & M \otimes_{\alpha_i \circ \alpha} S'' \\ & & & & \simeq \downarrow \theta & & \theta_2 \downarrow \theta_1 \\ 0 & \longrightarrow & N \simeq N \otimes_{\text{Id}_R} R & \xrightarrow{\text{Id}_N \otimes \alpha} & N \otimes_\alpha S & \begin{array}{c} \xrightarrow{\text{Id}_N \otimes \alpha_1} \\ \xrightarrow{\text{Id}_N \otimes \alpha_2} \end{array} & N \otimes_{\alpha_i \circ \alpha} S'' \end{array} \quad (2.5)$$

The rows are exact since $\alpha : R \rightarrow S$ is faithfully flat, e.g., $\text{Id}_M \otimes \alpha$ is injective and its image $M \otimes_{1_S}$ is the R -submodule $M \otimes_\alpha S$ where $\text{Id}_M \otimes \alpha_1$ and $\text{Id}_M \otimes \alpha_2$ agree.

The S/R -cocycle u defining the S/R -form N is

$$u = \theta_2^{-1} \circ \theta_1 \in \text{Aut}_{S''}(M \otimes_{\alpha_1 \circ \alpha} S'') = \text{Aut}_{S''}(M \otimes_R S'')$$

as we now explain.^b Let

$$L = \{x \in M \otimes_{\alpha} S : u((\text{Id}_M \otimes \alpha_1)(x)) = (\text{Id}_M \otimes \alpha_2)(x)\}. \quad (2.6)$$

It is clear that L is an R -submodule of $M \otimes_{\alpha} S = M \otimes_R S$, and a simple diagram chase in (2.5) above shows that the restriction of θ to L induces an isomorphism with $N \otimes 1_S \subset N \otimes_{\alpha} S = N \otimes_R S$. In other words, up to R -module isomorphism, our module N corresponds to the cocycle u .

It is well-known (and easy to check in any case) that u is a cocycle, i.e. that $u \otimes \alpha_{1,3} = (u \otimes \alpha_{2,3}) \circ (u \otimes \alpha_{1,2})$ where the $\alpha_{i,j} : S'' \rightarrow S''' = S \otimes_{\alpha} S \otimes_{\alpha} S$ are the natural S'' -algebra morphisms defined by putting 1_S in the position $l \neq i, j$. The cocycle condition can be rewritten in the form $\theta_{1,3} = \theta_{2,3} \circ \theta_{1,2}$ where the $\theta_{i,j}$ are automorphisms of the S''' -module $M \otimes_{\alpha_i, j \circ \alpha_i \circ \alpha} S''' = M \otimes_R S'''$ defined using a diagram similar to (2.4).

For R -algebras the situation is identical. Say that both A and B are R -algebras and that our isomorphism θ above is now an S -algebra isomorphism. Then u is an S'' -algebra automorphism of $A \otimes_R S''$, the descended R -module L is an R -subalgebra of $A \otimes_{\alpha} S = A \otimes_R S$ and the restriction of θ induces an R -algebra isomorphism between L and $B \simeq B \otimes 1_S$.

Theorem 2.9. *Let A be an R -algebra, and let B be an S/R form of A determined by the cocycle u as described in Descent Theory 2.8 above. If $F : k\text{-ALG} \rightarrow k\text{-MOD}$ is a functor over $k\text{-alg}$ stable under base change, then $F_R(B)$ is an S/R -form of the R -module $F_R(A)$ which is isomorphic as an R -module to the one given by the cocycle*

$$\nu_{A, \alpha_2 \circ \alpha}^{-1} \circ F(u) \circ \nu_{A, \alpha_1 \circ \alpha} \in \text{Aut}_{S''}(F_R(A) \otimes_R S'').$$

Proof. We fix an S -algebra isomorphism $\theta : A \otimes_{\alpha} S \rightarrow B \otimes_{\alpha} S$. The cocycle determining B (up to R -algebra isomorphism) is then $u = \theta_2^{-1} \circ \theta_1 \in \text{Aut}_{S''}(A \otimes_R S'')$. The result to establish can thus be rephrased by saying that

- (a) $z := (\nu_{A, \alpha_2 \circ \alpha}^{-1} \circ F(\theta_2)^{-1} \circ \nu_{B, \alpha_2 \circ \alpha}) \circ (\nu_{B, \alpha_1 \circ \alpha}^{-1} \circ F(\theta_1) \circ \nu_{A, \alpha_1 \circ \alpha}) \in \text{Aut}_{S''}(F_R(A) \otimes_R S'')$ is a cocycle.
- (b) The R -module determined by the cocycle z is isomorphic to $F_R(B)$.

^bNote that u can indeed be viewed as an S'' -module automorphism $M \otimes_{\alpha_1 \circ \alpha} S''$ because $\alpha_1 \circ \alpha = \alpha_2 \circ \alpha$. In what follows we will view S'' as an R -algebra via either one of these two (equal) maps. The notation $M \otimes_R S''$ responds to this convention.

Let us define $F^\nu(\theta) = \nu_{B,\alpha}^{-1} \circ F(\theta) \circ \nu_{A,\alpha}$ by means of the diagram

$$\begin{array}{ccc} F_R(A) \otimes_\alpha S & \xrightarrow{F^\nu(\theta)} & F_R(B) \otimes_\alpha S \\ \nu_{A,\alpha} \downarrow & & \uparrow (\nu_{B,\alpha})^{-1} \\ F_S(A \otimes_\alpha S) & \xrightarrow{F(\theta)} & F_S(B \otimes_\alpha S) \end{array}$$

In view of the descent of modules construction explained in Descent Theory 2.8, to prove (a) and (b) it will suffice to show that

$$\nu_{B,\alpha_i \circ \alpha}^{-1} \circ F(\theta_i) \circ \nu_{A,\alpha_i \circ \alpha} = (F^\nu(\theta))_i \quad \text{for } i = 1, 2. \quad (2.7)$$

Both cases $i = 1, 2$ are similar and we check in detail the case $i = 1$ only. We will use the following commutative diagram of S'' -module isomorphisms:

$$\begin{array}{ccc} F_R(A) \otimes_{\alpha_1 \circ \alpha} S'' & \xrightarrow{(F^\nu(\theta))_1} & F_R(B) \otimes_{\alpha_1 \circ \alpha} S'' \\ \downarrow (\text{Id}_{F_R(A)} \otimes \tau_1)^{-1} & & \downarrow (\text{Id}_{F_R(B)} \otimes \tau_1)^{-1} \\ F_R(A) \otimes_\alpha S \otimes_{\alpha_1} S'' & \xrightarrow{F^\nu(\theta) \otimes \text{Id}_{S''}} & F_R(B) \otimes_\alpha S \otimes_{\alpha_1} S'' \\ \downarrow \nu_{A,\alpha} \otimes \text{Id}_{S''} & & \downarrow \nu_{B,\alpha} \otimes \text{Id}_{S''} \\ F_S(A \otimes_\alpha S) \otimes_{\alpha_1} S'' & \xrightarrow{F(\theta) \otimes \text{Id}_{S''}} & F_S(B \otimes_\alpha S) \otimes_{\alpha_1} S'' \\ \downarrow \nu_{A \otimes_\alpha S, \alpha_1} & & \downarrow \nu_{B \otimes_\alpha S, \alpha_1} \\ F_{S''}((A \otimes_\alpha S) \otimes_{\alpha_1} S'') & \xrightarrow{F(\theta \otimes_{\alpha_1} \text{Id}_{S''})} & F_{S''}((B \otimes_\alpha S) \otimes_{\alpha_1} S'') \\ \downarrow F(\tau_1^A) & & \downarrow F(\tau_1^B) \\ F_{S''}(A \otimes_{\alpha_1 \circ \alpha} S) & \xrightarrow{F(\theta_1)} & F_{S''}(B \otimes_{\alpha_1 \circ \alpha} S'') \end{array}$$

The top rectangle commutes by definition of $F^\nu(\theta_1)$; the second rectangle commutes by applying the base change $\alpha_1 : S \rightarrow S''$ to the diagram defining $F^\nu(\theta)$; the third rectangle commutes by Lemma 2.4(a) for $(R, A), (S, B), (\alpha, f)$ and β replaced by $(S, A \otimes_\alpha S), (S, B \otimes_\alpha S), (\text{Id}_S, \theta)$ and $\text{Id}_{S''}$ respectively; the bottom rectangle commutes by applying F to the diagram (2.4) defining θ_1 . For the proof of (2.7) it is therefore sufficient to show that the dotted maps equal $\nu_{A,\alpha_1 \circ \alpha}$ and $\nu_{B,\alpha_1 \circ \alpha}$ respectively. We check the case of $\nu_{A,\alpha_1 \circ \alpha}$ by following explicitly the arrows on the left of the diagram. The case of $\nu_{B,\alpha_1 \circ \alpha}$, which is similar, is left to the reader. Let

$m \in F_R(A)$ and $s_1, s_2 \in S$. Then

$$\begin{aligned}
 m \otimes s_1 \otimes s_2 &\xrightarrow{(\text{Id}_{F_R(A)} \otimes \tau_1)^{-1}} m \otimes s_1 \otimes 1_S \otimes s_2 \\
 &\xrightarrow{\nu_{A, \alpha} \otimes \text{Id}_{S''}} (s_1 F_\alpha(\alpha^A)(m)) \otimes (1_S \otimes s_2) \\
 &\xrightarrow{\nu_{A \otimes_\alpha S, \alpha_1}} (1_S \otimes s_2)(F_{\alpha_1}(\alpha_1^{A \otimes_\alpha S})(s_1 F_\alpha(\alpha^A)(m))) \\
 &= (1_S \otimes s_2)(\alpha_1(s_1) F_{\alpha_1}(\alpha_1^{A \otimes_\alpha S})(F_\alpha(\alpha^A)(m))) \\
 &= (1_S \otimes s_2)(s_1 \otimes 1_S)(F_{\alpha_1}(\alpha_1^{A \otimes_\alpha S})(F_\alpha(\alpha^A)(m))) \\
 &= (s_1 \otimes s_2)(F_{\alpha_1}(\alpha_1^{A \otimes_\alpha S})(F_\alpha(\alpha^A)(m))) \\
 &\xrightarrow{F(\tau_1^A)} (s_1 \otimes s_2) F(\tau_1^A)(F_{\alpha_1}(\alpha_1^{A \otimes_\alpha S})(F_\alpha(\alpha^A)(m))) \\
 &= (s_1 \otimes s_2) F_{\alpha_1 \circ \alpha}((\alpha_1 \circ \alpha)^A(m)).
 \end{aligned}$$

This completes the proof since by definition $\nu_{A, \alpha_1 \circ \alpha}(m \otimes s_1 \otimes s_2) = (s_1 \otimes s_2) F_{\alpha_1 \circ \alpha}((\alpha_1 \circ \alpha)^A(m))$. \square

Remark 2.10. In the case that we are most interested in, the R -algebra A is of the form $A = \mathfrak{a} \otimes_k R$ for some k -algebra \mathfrak{a} . The isomorphism θ can now be thought as an S -algebra isomorphism (also denoted by θ)

$$\theta : \mathfrak{a} \otimes_k S \simeq (\mathfrak{a} \otimes_k R) \otimes_R S = A \otimes_R S \rightarrow B \otimes_R S,$$

where we have denoted \otimes_α by \otimes_R . Under the canonical S'' -isomorphism $\mathfrak{a} \otimes_k S'' \simeq (\mathfrak{a} \otimes_k R) \otimes_R S''$ (where we recall that S'' is viewed as an R -algebra via $\alpha_1 \circ \alpha = \alpha_2 \circ \alpha$) we can view the corresponding cocycle u as an S'' -algebra automorphism of $\mathfrak{a} \otimes_k S''$. The descended module L is then given by

$$L = \{x \in \mathfrak{a} \otimes_k S : u((\text{Id}_\mathfrak{a} \otimes \alpha_1)(x)) = (\text{Id}_\mathfrak{a} \otimes \alpha_2)(x)\}. \quad (2.8)$$

For future reference we record the following “descent setting” that covers the case that we are most interested in:

$$\left\{ \begin{array}{l}
 \mathfrak{a} \text{ is an algebra over } k; \\
 R \in k\text{-alg is such that } R/k \text{ is flat;} \\
 S \in R\text{-alg is such that } S/R \text{ is faithfully flat;} \\
 B \text{ is an } R\text{-algebra which is an } S/R\text{-form of } A = \mathfrak{a} \otimes_k R.
 \end{array} \right. \quad (2.9)$$

3. Invariant Functions

Unless stated otherwise, k is a commutative associative unital ring, R is a commutative associative unital k -algebra, namely $R \in k\text{-alg}$ in the notation of Sec. 2, M is an R -module, V is a k -module and B is an arbitrary (not necessarily associative, unital...) R -algebra, i.e. $(R, B) \in k\text{-ALG}$. Our goal is to study invariant bilinear functions $B \times B \rightarrow V$. We begin with pertinent definitions.

Definition 3.1 (((R, k) -bilinear functions). A k -bilinear function $\beta : M \times M \rightarrow V$ is called (R, k) -bilinear if $\beta(rm_1, m_2) = \beta(m_1, rm_2)$ holds for all $m_i \in M$ and $r \in R$.

We denote by $\mathcal{L}_{(R,k)}^2(M; V)$ the R -module of (R, k) -bilinear maps $M \times M \rightarrow V$. Its R -module structure is given by $(r\beta)(m_1, m_2) = \beta(rm_1, m_2)$ for $r \in R$ and $\beta \in \mathcal{L}_{(R,k)}^2(M; V)$. We abbreviate $\mathcal{L}_R^2(M) = \mathcal{L}_{(R,R)}^2(M; R)$.

It is immediate from [8, II, Sec. 4.1] that one has a commutative triangle

$$\begin{array}{ccc}
 \text{Hom}_k(M \otimes_R M, V) & \xrightarrow[\simeq]{\times} & \mathcal{L}_{(R,k)}^2(M; V) \\
 & \searrow[\simeq]{y} & \nearrow[\simeq]{z} \\
 & & \text{Hom}_R(M, \text{Hom}_k(M, V))
 \end{array} \tag{3.1}$$

of R -linear isomorphisms: For $\varphi \in \text{Hom}_k(M \otimes_R M, V)$ one defines $(\times(\varphi))(m_1, m_2) = \varphi(m_1 \otimes m_2)$. This is an R -linear map with respect to the R -action on $\text{Hom}_k(M \otimes_R M, V)$ given by $(r\varphi)(m_1 \otimes m_2) = \varphi(rm_1 \otimes m_2)$ for $r \in R$. Also, $(y(\varphi))(m_1)$ maps m_2 onto $\varphi(m_1 \otimes m_2)$. The map y is R -linear if we view $\text{Hom}_k(M, V)$ as an R -module via $(rh)(m) = h(rm)$ for $h \in \text{Hom}_k(M, V)$. To $\beta \in \text{Hom}_R(M, \text{Hom}_k(M, V))$ one associates the (R, k) -bilinear function $z(\beta)$ defined by $(z(\beta))(m_1, m_2) = (\beta(m_1))(m_2)$.

One calls $\beta \in \mathcal{L}_{(R,k)}^2(M; k)$ *nondegenerate* (respectively, *nonsingular*) if $z^{-1}(\beta) \in \text{Hom}_R(M, M^*)$, $M^* = \text{Hom}_k(M, k)$, is injective (respectively, bijective).^c In more familiar terms $\beta \in \mathcal{L}_{(R,k)}^2(M; k)$ is nondegenerate if and only if $\beta(b, M) = 0$ implies $b = 0$. If k is a field, the existence of a nonsingular bilinear form on a k -algebra B forces B to be of finite dimension. Moreover, for a finite-dimensional B a form is nondegenerate if and only if it is nonsingular. See Lemma 6.16 for a generalization.

Definition 3.2 (Invariant functions). We call $\beta \in \mathcal{L}_{(R,k)}^2(B; V)$ *invariant* if

$$\beta(ab, c) = \beta(a, bc) = \beta(b, ca) \tag{3.2}$$

holds for all $a, b, c \in B$.

Clearly, the set $\text{IBF}_{(R,k)}(B; V)$ of all invariant (R, k) -bilinear functions $B \times B \rightarrow V$ is a submodule of the R -module $\mathcal{L}_{(R,k)}^2(B; V)$. The following special cases of $\text{IBF}_{(R,k)}(B; V)$ are of particular interest:

$$\begin{aligned}
 \text{IBF}_k(B; V) &:= \text{IBF}_{(k,k)}(B; V), \\
 \text{IBF}_{(R,k)}(B) &:= \text{IBF}_{(R,k)}(B; k), \\
 \text{IBF}_k(B) &:= \text{IBF}_k(B; k) = \text{IBF}_{(k,k)}(B).
 \end{aligned}$$

The elements of $\text{IBF}_k(B)$ are called *invariant k -bilinear forms*. Of particular importance is the case $k = R$; these are the invariant R -bilinear forms on B .

Remark 3.3. The above definition works equally well for invariant bilinear functions on dimodules of algebras (for the definition of a dimodule see [23] as well

^cThe asymmetry in these definitions (one should, strictly speaking, speak of left and right nondegeneracy and nonsingularity) will not play a major role in this paper since our main interest later will be in invariant bilinear forms of perfect algebras which, by Remark 3.4, are symmetric.

as Lemma 3.7 supra). This is not without interest as these types of bilinear forms are an important tool for the study of the representation theory of the algebras in question, for example for the existence of the Jantzen filtration of Verma modules. We will not pursue this more general set up in this work.

Remark 3.4. If B is perfect, namely if $B = BB$, where $BB = \text{Span}_R\{ab : a, b, \in B\}$, every invariant bilinear function is symmetric: $\beta(ab, c) = \beta(a, bc) = \beta(b, ca) = \beta(bc, a) = \beta(c, ab)$. Moreover, any k -bilinear function is already (R, k) -bilinear:

$$\text{IBF}_k(B; V) = \text{IBF}_{(R,k)}(B; V) \quad (B \text{ perfect}).$$

Indeed, $\beta(r(ab), c) = \beta(a(rb), c) = \beta(a, (rb)c) = \beta(a, b(rc)) = \beta(ab, rc)$.

Definition 3.5 (Universal invariant function). Let $\mathbf{IBF}_R(B)$ be the quotient of the R -module $B \otimes_R B$ by the submodule

$$\begin{aligned} \text{ibf}_R(B) &= \text{Span}_R\{ab \otimes c - a \otimes bc, ab \otimes c - b \otimes ca : a, b, c \in B\} \\ &= \text{Span}_{\mathbb{Z}}\{ab \otimes c - a \otimes bc, ab \otimes c - b \otimes ca : a, b, c \in B\} \\ &= \text{Span}_{\mathbb{Z}}\{ab \otimes c - a \otimes bc, a \otimes bc - bc \otimes a : a, b, c \in B\}, \end{aligned} \quad (3.3)$$

where the last equality follows from $(bc \otimes a - a \otimes bc) + (ab \otimes c - b \otimes ca) = (ab \otimes c - a \otimes bc) + (bc \otimes a - b \otimes ca)$. Denoting by $i_B : \text{ibf}_R(B) \rightarrow B \otimes_R B$ the inclusion and by q_B the canonical quotient map

$$q_B : B \otimes_R B \rightarrow \mathbf{IBF}_R(B), \quad a \otimes b \mapsto \overline{a \otimes b},$$

we have an exact sequence of R -modules

$$0 \rightarrow \text{ibf}_R(B) \xrightarrow{i_B} B \otimes_R B \xrightarrow{q_B} \mathbf{IBF}_R(B) \rightarrow 0. \quad (3.4)$$

We define an invariant R -bilinear function

$$\beta_{\text{uni}} : B \times B \rightarrow \mathbf{IBF}_R(B), \quad \beta_{\text{uni}}(a, b) = \overline{a \otimes b}, \quad (3.5)$$

called the *universal invariant R -bilinear function*.^d This terminology is justified because of the following natural R -module isomorphism

$$\text{Hom}_k(\mathbf{IBF}_R(B), V) \xrightarrow{\cong} \text{IBF}_{(R,k)}(B; V), \quad f \mapsto \tilde{f} = f \circ \beta_{\text{uni}}. \quad (3.6)$$

Its inverse

$$\text{IBF}_{(R,k)}(B; V) \xrightarrow{\cong} \text{Hom}_k(\mathbf{IBF}_R(B), V), \quad \beta \mapsto \bar{\beta} \quad (3.7)$$

assigns to $\beta \in \text{IBF}_{(R,k)}(B, V)$ the unique k -linear map

$$\bar{\beta} : \mathbf{IBF}_R(B) \rightarrow V, \quad \bar{\beta}(\overline{a \otimes b}) = \beta(a, b). \quad (3.8)$$

In other words, $\mathbf{IBF}_R(B)$ represents the obvious functor $\text{IBF}(B; -) : k\text{-mod} \rightarrow R\text{-mod}$. The isomorphism (3.6) determines $\mathbf{IBF}_R(B)$ up to a unique k -linear isomorphism. We will describe $\mathbf{IBF}_R(B)$ for several cases of interest in this paper.

^dWe want to thank K.-H. Neeb for bringing this concept to our attention.

The most important situation is captured by the following result that we state in the form of a principle.

IBF-principle 3.6. *Assume B is an R -algebra for some $R \in k\text{-alg}$. Let $\beta \in \text{IBF}_R(B)$ be such that the induced map $\bar{\beta} : \mathbf{IBF}_R(B) \rightarrow R, \overline{b_1 \otimes b_2} \mapsto \beta(b_1, b_2)$ is an R -module isomorphism. Then for any k -module V the map*

$$\text{Hom}_k(R, V) \rightarrow \text{IBF}_{(R,k)}(B; V), \quad \varphi \mapsto \varphi \circ \beta \tag{3.9}$$

is an isomorphism of R -modules. In particular,

- (a) *the map $R^* = \text{Hom}_k(R, k) \rightarrow \text{IBF}_{(R,k)}(B), \varphi \mapsto \varphi \circ \beta$ is an isomorphism, and*
- (b) *every $\gamma \in \text{IBF}_R(B)$ is of the form $\gamma = r\beta$ for a unique $r \in R$.*

Proof. This follows from the isomorphism (3.6) and the equality $\beta = \bar{\beta} \circ \beta_{\text{uni}}$. □

We will say that (B, β) satisfies the *IBF-principle* if the assumptions and hence also the conclusions of IBF-principle 3.6 hold. Note that in this case we have a precise and explicit description of *all* invariant (R, k) -bilinear functions on B .

While the connection between invariant bilinear forms and centroids does not feature prominently in this paper, it has nevertheless been an important guiding principle for our work. Besides the conceptual importance of this connection, another reason for elaborating on it is Corollary 3.7 relating invariant forms and the centroid of an algebra. Not only will this provide the reader with a means to determine the module $\text{IBF}_{(R,k)}(B)$, but it will also be useful in Sec. 6 when we will be looking at algebras with a “one-dimensional” $\text{IBF}_{(R,k)}(B)$.

In preparation for these results, we first present the necessary background. Using the terminology of [23], we recall that a (B, R) -*dimodule* is an R -module M together with R -bilinear maps $B \times M \rightarrow M$ and $M \times B \rightarrow M$. For example, B itself is a (B, R) -dimodule with respect to the left and right multiplications of the algebra B , called the *regular dimodule* and denoted B_{reg} . Also the R -module $\text{Hom}_k(B, V)$ is a (B, R) -dimodule with respect to the B -actions $(b_1 \cdot \varphi)(b_2) = \varphi(b_2 b_1) = (\varphi \cdot b_2)(b_1)$. For any (B, R) -dimodule M the *centroid of B with values in M* is

$$\begin{aligned} \text{Ctd}_R(B, M) &= \{ \chi \in \text{Hom}_R(B, M) : \chi(b_1 b_2) = b_1 \cdot \chi(b_2) = \chi(b_1) \cdot b_2 \\ &\text{for all } b_1, b_2 \in B \}. \end{aligned}$$

Taking as M the regular dimodule B_{reg} , we recover the usual notion of the centroid of B : $\text{Ctd}_R(B) = \text{Ctd}_R(B, B_{\text{reg}})$. We note that $\text{Ctd}_R(B)$ is isomorphic to the usual center if B is a unital algebra, where, we recall, the center of an arbitrary algebra B consists of those $c \in B$ which commute with all $b \in B$, i.e. $cb = bc$, and which associate with all $b_1, b_2 \in B$, i.e. $(c, b_1, b_2) = (b_1, c, b_2) = (b_1, b_2, c) = 0$ where $(x, y, z) = (xy)z - x(yz)$. There exists a natural map $R \rightarrow \text{Ctd}_R(B)$ given by

$r \mapsto \chi_r$ where $\chi_r(b) = rb$ for all $b \in B$. We call B a *central R -algebra* if this map is an isomorphism. In this situation we will often identify R with $\text{Ctd}_R(B)$ without any further explicit reference.

We can now describe the connection between invariant functions and centroids.

Lemma 3.7. *The restriction of the R -isomorphism z of (3.1) to $\text{Ctd}_R(B, \text{Hom}_k(B, V))$ induces an R -linear isomorphism $\text{Ctd}_R(B, \text{Hom}_k(B, V)) \simeq \text{IBF}_{(R,k)}(B; V)$. In particular,*

$$\text{Ctd}_R(B, B^*) \simeq \text{IBF}_{(R,k)}(B).$$

Proof. The result is a straightforward consequence of the various definitions. \square

Corollary 3.8. *Assume $\beta \in \text{IBF}_{(R,k)}(B)$ is nonsingular. Then*

$$\text{Ctd}_R(B) \simeq \text{IBF}_{(R,k)}(B) \quad (\text{isomorphism of } R\text{-modules}).$$

If furthermore B is a central R -algebra, then $\text{IBF}_{(R,k)}(B)$ is a free R -module of rank 1 admitting β as a basis:

$$\text{IBF}_{(R,k)}(B) = R\beta \simeq R. \tag{3.10}$$

Proof. By assumption $\chi_\beta = z^{-1}(\beta) \in \text{Ctd}_R(B, B^*)$ is an isomorphism of R -modules. The fact that χ_β is centroidal amounts to saying that $B_{\text{reg}} \simeq B^*$ as dimodules, whence $\text{IBF}_{(R,k)}(B) \simeq \text{Ctd}_R(B, B^*) \simeq \text{Ctd}_R(B) \simeq R$. Via these isomorphism we have $\beta \mapsto \chi_\beta \mapsto \chi_\beta^{-1} \circ \chi_\beta = \text{Id}_B \mapsto 1_R$. Since 1_R is a basis of R , β is a basis of $\text{IBF}_{(R,k)}(B)$. \square

As we will see in Sec. 6, there are many natural types of algebras satisfying the assumptions of Corollary 3.8. The following result will allow us to transition from the general setting to the specific examples.

Proposition 3.9. *Assume that B is a central R -algebra, that $\beta \in \text{IBF}_R(B)$ is nonsingular with $\beta(B, B) = R$, and that $\mathbf{IBF}_R(B)$ is projective. Then the IBF-principle 3.6 holds for (B, β) .*

Proof. By assumption $\bar{\beta}$ is surjective. Denoting by K the kernel of $\bar{\beta}$, we obtain a split exact sequence

$$0 \rightarrow K \rightarrow \mathbf{IBF}_R(B) \xrightarrow{\bar{\beta}} R \rightarrow 0$$

of R -modules and consequently a split exact sequence

$$0 \rightarrow \text{Hom}_R(R, R) \xrightarrow{\hat{\beta}} \text{Hom}_R(\mathbf{IBF}_R(B), R) \rightarrow \text{Hom}_R(K, R) \rightarrow 0$$

with $\hat{\beta}(r \text{Id}_R) = r \text{Id}_R \circ \bar{\beta} = r\bar{\beta}$ for $r \in R$. The R -module isomorphism (3.6) sends $\bar{\beta}$ to β . We also know that $\text{IBF}_R(B) \simeq R$ by Corollary 3.8 (applied to the case

$k = R$). Hence the diagram

$$\begin{array}{ccc}
 \mathrm{Hom}_R(R, R) & \xrightarrow{\hat{\beta}} & \mathrm{Hom}_R(\mathbf{IBF}_R(B), R) & & r \mathrm{Id}_R & \xrightarrow{\quad} & r\bar{\beta} \\
 \uparrow \simeq & & \downarrow \simeq & & \uparrow & & \downarrow \\
 R & \xrightarrow{\simeq} & \mathrm{IBF}_R(B) & & r & \xrightarrow{\quad} & r\beta
 \end{array}$$

commutes, proving that $\hat{\beta}$ is an isomorphism. This in turn forces $K^{(*)} := \mathrm{Hom}_R(K, R) = \{0\}$.^e Since $\mathbf{IBF}_R(B) \simeq K \oplus R$ is projective, so is K . But then the canonical map $K \rightarrow (K^{(*)})^{(*)} = 0$ is injective. Thus $K = 0$. \square

4. Functorial Nature of IBF

As in the previous section k will denote a commutative associative unital ring and $R \in k\text{-alg}$. The main purpose of this section is to describe the functorial nature of \mathbf{IBF} and study its behavior under base change.

Let $(\alpha, f): (R, M) \rightarrow (S, N)$ be a morphism in $k\text{-MOD}$ (see Definition 2.1), i.e. $\alpha: R \rightarrow S$ is a morphism in $k\text{-alg}$ and $f: M \rightarrow N$ is α -semilinear, and suppose V is a k -module. Then for any $\kappa \in \mathcal{L}_{(S,k)}^2(N; V)$ the map

$$f^*(\kappa): M \times M \rightarrow V, \quad (m_1, m_2) \mapsto \kappa(f(m_1), f(m_2)) \quad (4.1)$$

is (R, k) -bilinear. We obtain in this way a k -linear map

$$f^*: \mathcal{L}_{(S,k)}^2(N; V) \rightarrow \mathcal{L}_{(R,k)}^2(M; V), \quad \kappa \mapsto f^*(\kappa).$$

If $(\beta, g): (S, N) \rightarrow (T, P)$ is another morphism in $k\text{-MOD}$, it is immediate that $(g \circ f)^* = f^* \circ g^*$. Observe that this in particular defines a right action of the group $\mathrm{GL}_R(M)$ on $\mathcal{L}_{(R,k)}^2(M; V)$ (see Remark 5.2 for a functorial version of this observation).

Base Change 4.1. Let $\kappa: M \times M \rightarrow R$ be an R -bilinear form and let $\alpha: R \rightarrow S$ be a morphism in $k\text{-alg}$. There exists a unique S -bilinear form $\kappa_\alpha: M \otimes_\alpha S \times M \otimes_\alpha S \rightarrow S$ satisfying $\kappa_\alpha(m_1 \otimes s_1, m_2 \otimes s_2) = \alpha(\kappa(m_1, m_2))s_1s_2$. In case α is clear from the context, we will denote $\kappa_\alpha = \kappa_S$ and call κ_S the *base change of κ by S* . We then have the equation $\kappa_S(m_1 \otimes s_1, m_2 \otimes s_2) = \kappa(m_1, m_2)s_1s_2$.

Base change can also be understood in terms of the isomorphism $\times: \mathrm{Hom}_R(M \otimes_R M, R) \xrightarrow{\simeq} \mathcal{L}_R^2(M)$ of (3.1). Let $\times^{-1}(\kappa) = \tilde{\kappa}: M \otimes_R M \rightarrow R$ be the R -linear map associated to κ . Then $\times^{-1}(\kappa_S) = \tilde{\kappa}_S$ is obtained from $\tilde{\kappa} \otimes \mathrm{Id}_S$ with the aid of two canonical S -module isomorphisms, namely

$$\tilde{\kappa}_S: (M \otimes_R S) \otimes_S (M \otimes_R S) \simeq (M \otimes_R M) \otimes_R S \xrightarrow{\tilde{\kappa} \otimes \mathrm{Id}_S} R \otimes_R S \simeq S.$$

The following lemma collects some results using base change.

^eTo avoid any possible confusion we use $(^*)$ as opposed to $*$ to denote the R -dual given that, by convention, $*$ always refers to the k -dual.

Lemma 4.2. (a) (Transitivity of base change) Let $\kappa \in \mathcal{L}_R^2(M)$, let $R \xrightarrow{\alpha} S \xrightarrow{\beta} T$ be morphisms in $k\text{-alg}$, and let $\zeta : (M \otimes_{\alpha} S) \otimes_{\beta} T \rightarrow M \otimes_{\beta \circ \alpha} T$, $m \otimes s \otimes t \mapsto m \otimes (\beta(s))t$, be the canonical isomorphism of T -modules. Then

$$\zeta^* \kappa_{\beta \circ \alpha} = (\kappa_{\alpha})_{\beta}. \quad (4.2)$$

(b) Let M and N be R -modules, $\lambda \in \mathcal{L}_R^2(N)$, $f : M \rightarrow N$ be an R -linear map and $\alpha : R \rightarrow S$ be a morphism in $k\text{-alg}$. Then

$$(f^*(\lambda))_{\alpha} = (f \otimes \text{Id}_S)^*(\lambda_{\alpha}). \quad (4.3)$$

(c) Let $\kappa, \kappa' \in \mathcal{L}_R^2(M)$. Then $\kappa = \kappa'$ if and only if $\kappa_S = \kappa'_S$ for some faithfully flat extension $S \in R\text{-alg}$.

(d) Assume that M is finitely presented. Let $\kappa \in \mathcal{L}_R^2(M)$ and let $S \in R\text{-alg}$ be such that S is a flat R -module. If κ is nondegenerate (respectively, nonsingular), then κ_S is nondegenerate (respectively, nonsingular). The converse holds in both cases if S/R is faithfully flat.

Proof. The proofs of (a) and (b) are immediate from the definitions. In (c) suppose $\kappa_S = \kappa'_S$ for some faithfully flat $S \in R\text{-alg}$. Then $(x^{-1}(\kappa))_S = x^{-1}(\kappa_S) = x^{-1}(\kappa'_S) = (x^{-1}(\kappa'))_S$ as maps $(M \otimes_R S) \otimes_S (M \otimes_R S) \rightarrow S$ by Base Change 4.1. But then $x^{-1}(\kappa) = x^{-1}(\kappa')$ by faithfully flat descent, whence $\kappa = \kappa'$.

(d) Since M is finitely presented and S/R is flat, the canonical map $\omega : \text{Hom}_R(M, R) \otimes_R S \rightarrow \text{Hom}_S(M \otimes_R S, S)$ is an isomorphism. Recall from (3.1) the R -linear map $z^{-1}(\kappa) : M \rightarrow \text{Hom}_R(M, R)$. It is immediate from the definitions that

$$\begin{array}{ccc} M \otimes_R S & \xrightarrow{z^{-1}(\kappa) \otimes \text{Id}_S} & \text{Hom}_R(M, R) \otimes_R S \\ & \searrow z^{-1}(\kappa_S) & \swarrow \omega \\ & \text{Hom}_S(M \otimes_R S, S) & \end{array}$$

is a commutative diagram. Hence $z^{-1}(\kappa_S)$ is injective (respectively, bijective) if and only if $z^{-1}(\kappa)$ is so. The claim then follows from standard properties of flat (respectively, faithfully flat) extensions. \square

Proposition 4.3 (IBF and ibf as functors). (a) Let $(\alpha, f) : (R, B) \rightarrow (S, C)$ be a morphism in $k\text{-ALG}$. The map

$$f \otimes f : B \otimes_R B \rightarrow C \otimes_S C, \quad b_1 \otimes b_2 \mapsto f(b_1) \otimes f(b_2)$$

is α -semilinear and maps $\text{ibf}_R(B)$ to $\text{ibf}_S(C)$. We denote by $\text{ibf}_{\alpha}(f) : \text{ibf}_R(B) \rightarrow \text{ibf}_S(C)$ the restricted map and by $\mathbf{IBF}_{\alpha}(f) : \mathbf{IBF}_R(B) \rightarrow \mathbf{IBF}_S(C)$ the induced quotient map.

(b) *The assignments*

$$(R, B) \rightarrow (R, \mathbf{IBF}_R(B)) \quad \text{and} \quad (\alpha, f) \mapsto (\alpha, \mathbf{IBF}_\alpha(f))$$

define a functor $\mathbf{IBF} : k\text{-ALG} \rightarrow k\text{-MOD}$ over $k\text{-alg}$ which is stable under base change in the sense of Definition 2.5.

(c) *The assignments*

$$(R, B) \rightarrow (R, \text{ibf}_R(B)) \quad \text{and} \quad (\alpha, f) \mapsto (\alpha, \text{ibf}_\alpha(f))$$

define a functor $\text{ibf} : k\text{-ALG} \rightarrow k\text{-MOD}$ over $k\text{-alg}$ which is stable under flat base change.

Proof. Part (a) is straightforward. We will prove (b) and (c) at the same time. It is easy to verify that \mathbf{IBF} and ibf are functors over $k\text{-alg}$.

We have already noted in Example 2.6 that $B \mapsto B \otimes_R B$ and $(\alpha, f) \mapsto (\alpha, f \otimes f)$ defines a functor over $k\text{-alg}$ which is stable under base change. Indeed, for any morphism $\alpha : R \rightarrow S$ the map $\nu_{B,\alpha}$ of (2.3) is the well-known isomorphism

$$\nu_{B,\alpha} : (B \otimes_R B) \otimes_R S \rightarrow (B \otimes_R S) \otimes_S (B \otimes_R S), \quad b_1 \otimes b_2 \otimes s \mapsto b_1 \otimes 1_R \otimes b_2 \otimes s.$$

In the following we will abbreviate $\nu = \nu_{B,\alpha}$ and $\bar{\nu} = \bar{\nu}_{B,\alpha} : \mathbf{IBF}_R(B) \otimes_\alpha S \rightarrow \mathbf{IBF}_S(B \otimes_\alpha S)$. We have the following diagram with exact rows

$$\begin{array}{ccccccc}
 0 & \xrightarrow{\text{(c)}} & \text{ibf}_R(B) \otimes_R S & \xrightarrow{i_B \otimes \text{Id}_S} & (B \otimes_R B) \otimes_R S & \xrightarrow{q_B \otimes \text{Id}_S} & \mathbf{IBF}_R(B) \otimes_R S \longrightarrow 0 \\
 & & \downarrow & & \simeq \downarrow \nu & & \downarrow \bar{\nu} \\
 0 & \longrightarrow & \text{ibf}_S(B \otimes_R S) & \xrightarrow{i_B \otimes_R S} & (B \otimes_R S) \otimes_S (B \otimes_R S) & \xrightarrow{q_B \otimes_R S} & \mathbf{IBF}_S(B \otimes_R S) \longrightarrow 0
 \end{array} \tag{4.4}$$

where the top row is obtained by tensoring (3.4) with S and the bottom row is (3.4) for $B \otimes_R S$ (under assumption (c), namely when S/R is flat, then $i_B \otimes \text{Id}_S$ is injective. This is reflected by the dashed line at the top left of the diagram). One easily verifies that ν sends the S -submodule $(i_B \otimes \text{Id}_S)(\text{ibf}_R(B) \otimes_R S)$ of $(B \otimes_R B) \otimes S$ onto $\text{ibf}_S(B \otimes_R S) \subset (B \otimes_R S) \otimes_S (B \otimes_R S)$. It follows that by restriction ν induces an S -module isomorphism between $(i_B \otimes \text{Id}_S)(\text{ibf}_R(B) \otimes_R S)$ and $\text{ibf}_S(B \otimes_R S)$. Since from the definitions of ν and $\bar{\nu}$ the right-hand side square of diagram (4.4) commutes, a simple diagram chase shows that $\bar{\nu}$ is an isomorphism. In case S is a flat extension, the dashed vertical arrow is injective, hence bijective. \square

Corollary 4.4. *Let B be an S/R -form of $A = \mathfrak{a} \otimes_k R$ as in the descent setting (2.9). Let $u \in \mathbf{Aut}(\mathfrak{a})(S'')$ be a cocycle determining B up to R -algebra isomorphism (see Remark 2.10). We denote by $\nu : \mathbf{IBF}_k(\mathfrak{a}) \otimes_k S'' \rightarrow \mathbf{IBF}_{S''}(\mathfrak{a} \otimes_k S'')$ the isomorphism (2.3) for $F = \mathbf{IBF}$. Then $\mathbf{IBF}_R(B)$ is an S/R -form of $\mathbf{IBF}(\mathfrak{a} \otimes_k R)$ which is isomorphic as an R -module to the one given by the cocycle $\nu^{-1} \circ \mathbf{IBF}(u) \circ \nu$.*

Proof. This is a special case of Theorem 2.9. \square

Remark 4.5. In the above corollary we can take $k = R$. The result then applies to an arbitrary R -algebra A .

Lemma 4.6. *Let B be an R -algebra, $\beta \in \mathbf{IBF}_R(B)$, and $S \in R\text{-alg}$.*

- (a) *We denote by $\nu: \mathbf{IBF}_R(B) \otimes_R S \rightarrow \mathbf{IBF}_S(B \otimes_R S)$ the isomorphism (2.3) for $F = \mathbf{IBF}$, by $(\bar{\beta})_S$ the base change of $\bar{\beta}: \mathbf{IBF}_R(B) \rightarrow R$ and by $(\overline{\beta_S})$ the map (3.8) associated to the bilinear form β_S . Then, after identifying $R \otimes_R S = S$, we have $(\bar{\beta})_S = \overline{(\beta_S)} \circ \nu$:*

$$\begin{array}{ccc} \mathbf{IBF}_R(B) \otimes_R S & \xrightarrow{(\bar{\beta})_S} & R \otimes_R S \\ \nu \downarrow \simeq & & \parallel \\ \mathbf{IBF}_S(B \otimes_R S) & \xrightarrow{\overline{(\beta_S)}} & S \end{array}$$

- (b) *Assume that B is finitely presented as an R -module and that S/R is flat. If (B, β) satisfies the IBF-principle 3.6, then so does (B_S, β_S) . The converse is true whenever S/R is faithfully flat.*

Proof. Part (a) is immediate from the definitions. In (b) we know from (a) that $\overline{\beta_S}$ is an isomorphism. The assertion about faithful flatness is standard. \square

5. Descent of Bilinear Forms

In this section we will study the descent of bilinear forms in the setting of (2.9). We are interested in having a full understanding of all k -bilinear forms on B . The guiding principle is that this ought to be possible if one knows the nature of the k -bilinear forms of \mathfrak{a} , for example if \mathfrak{a} is a finite-dimensional central-simple Lie algebra over a field of characteristic 0 (Theorem 6.5).

That this may be possible at all, is somehow surprising. The twisted nature and descent theory related to B view B as an object over R (note that it is not the case that B is in any meaningful way a twisted form of $\mathfrak{a} \otimes_k R$ or \mathfrak{a} as k -algebras). Yet the information that we will get is about k -bilinear forms $B \times B \rightarrow k$. As mentioned before, it is the k - (and not R -) bilinear forms on B which are often of interest (such as in the case of infinite-dimensional Lie theory as exemplified, for example, by the affine Kac–Moody Lie algebras. See also Secs. 6 and 7 below).

Assume that κ is a k -bilinear form on \mathfrak{a} . We want to know when, in a natural fashion, we can attach to κ an R -bilinear form κ_B on B . It will be κ_B that will lead us to fully understand all k -bilinear functions on B . The key assumption that makes this construction possible and natural is that κ be invariant under algebra automorphisms. We define this concept before proceeding with the main results.

Automorphism Invariance 5.1. For an R -algebra B we denote by $\mathbf{Aut}(B)$ the automorphism group functor of B . We remind the reader that $\mathbf{Aut}(B)$ is the functor from the category $R\text{-alg}$ to the category of groups that attaches to $S \in R\text{-alg}$ the

group $\text{Aut}_S(B \otimes_R S)$ of automorphisms of the S -algebra $B \otimes_R S$, and to an arrow $S \rightarrow T$ in $R\text{-alg}$ and $f \in \text{Aut}_S(B \otimes_R S)$ the automorphism $f \otimes \text{Id}_T$ of $B \otimes_R T \simeq (B \otimes_R S) \otimes_S T$. We say that $\beta \in \mathcal{L}_R^2(B)$ is **Aut**(B)-invariant if $f^*(\beta_S) = \beta_S$ holds for all $S \in R\text{-alg}$ and all $f \in \mathbf{Aut}(B)(S)$, where we remind the reader that $f^*(\beta_S)(b_1 \otimes s_1, b_2 \otimes s_2) = \beta_S(f(b_1 \otimes s_1), f(b_2 \otimes s_2))$. In other words, β is **Aut**(B)-invariant if and only if β_S is $\text{Aut}_S(B \otimes_R S)$ -invariant in the obvious sense for all $S \in R\text{-alg}$.

Remark 5.2. Automorphism invariance has a functorial interpretation. Namely, we have a functor $\mathcal{L}^2(B) : R\text{-alg} \rightarrow R\text{-MOD}$ which assigns to the extension S/R the S -module $\mathcal{L}_S^2(B \otimes_R S)$, and is given at the level of arrows by Base Change 4.1. The automorphism group functor **Aut**(B) acts on the functor $\mathcal{L}^2(B)$ from the right. A bilinear form $\beta \in \mathcal{L}_R^2(B)$ is **Aut**(B)-invariant if and only if it is invariant under this action.

In particular, the above considerations apply to modules, viewed as trivial algebras. If M is an R -module, the R -group functor $\mathbf{GL}(M)$ acts naturally on $\mathcal{L}_R^2(M)$.

Automorphism invariance behaves nicely with respect to base change and faithfully flat descent.

Lemma 5.3. *Let B be an R -algebra, $\beta \in \mathcal{L}_R^2(B)$ and $S \in R\text{-alg}$. If β is **Aut**(B)-invariant, then β_S is **Aut**($B \otimes_R S$)-invariant. The converse holds if S/R is faithfully flat.*

Proof. (I) We begin with a general observation. Let T be an extension of S . The canonical T -linear algebra isomorphism $\zeta : (B \otimes_R S) \otimes_S T \rightarrow B \otimes_R T$ of Lemma 4.2(a) induces a group isomorphism

$$\text{Aut}_T((B \otimes_R S) \otimes_S T) \xrightarrow{\simeq} \text{Aut}_T(B \otimes_R T), \quad f \mapsto \zeta \circ f \circ \zeta^{-1}$$

(we view T as an object in $R\text{-alg}$ in the obvious way). Since $(\zeta^{-1})^*((\beta_S)_T) = \beta_T$ by (4.2), it follows that $(\beta_S)_T$ is $\text{Aut}_T((B \otimes_R S) \otimes_S T)$ -invariant if and only if β_T is $\text{Aut}_T(B \otimes_R T)$ -invariant.

(II) It is immediate from (I) and the definitions that if β is **Aut**(B)-invariant then β_S is **Aut**($B \otimes S$)-invariant.

(III) Assume now that S/R is faithfully flat and that β_S is **Aut**($B \otimes_R S$)-invariant. To prove that β is **Aut**(B)-invariant, let $S' \in R\text{-alg}$ and let $f \in \mathbf{Aut}(B)(S')$. The extension $T = S \otimes_R S'$ of S' is faithfully flat. Hence, by Lemma 4.2(c), we have $f^*(\beta_{S'}) = \beta_{S'}$ as soon as $(f^*(\beta_{S'}))_T = (\beta_{S'})_T \in \mathcal{L}_T^2((B \otimes_R S') \otimes_{S'} T)$. Note that $(f^*(\beta_{S'}))_T = (f \otimes \text{Id}_T)^*((\beta_{S'})_T)$ by (4.3). Applying the considerations of (I) to the isomorphism $\zeta' : (B \otimes_R S') \otimes_{S'} T \rightarrow B \otimes_R T$ shows that we need to prove that β_T is $\text{Aut}_T(B \otimes_R T)$ -invariant. But by (I) again this is indeed the case. □

Theorem 5.4 (Descent of Aut-invariant forms). *Assume that we are under the descent setting of (2.9): \mathfrak{a} is a k -algebra, $R \in k\text{-alg}$ is flat, and we are given an*

R -algebra B which is a twisted form of $A = \mathfrak{a} \otimes_k R$, hence split by some faithfully flat extension of R . Assume that $\kappa \in \mathcal{L}_k^2(\mathfrak{a})$ is an $\mathbf{Aut}(\mathfrak{a})$ -invariant bilinear form.

- (a) There exists a unique R -bilinear form $\kappa_B \in \mathcal{L}_R^2(B)$ such that $(\kappa_B)_S = \theta^*(\kappa_S)$ whenever S/R is faithfully flat and $\theta: B \otimes_R S \rightarrow \mathfrak{a} \otimes_k S$ is an isomorphism of S -algebras. Moreover, κ_B is $\mathbf{Aut}(B)$ -invariant.
- (b) If κ is invariant, then so is κ_B .
- (c) If \mathfrak{a} is finitely presented and κ is nondegenerate (respectively, nonsingular), then so is κ_B .

Proof. (a) Throughout the proof we fix S and θ as in (a). We let $\alpha: R \rightarrow S$ be the structure map and κ_S the base change of κ to S . It will be convenient to first point out a general observation, which is immediate from the definitions.

$$\text{If } \beta: S \rightarrow T \text{ is a homomorphism in } k\text{-alg}, \text{ then } (\text{Id}_{\mathfrak{a}} \otimes \beta)^*(\kappa_T) = \beta \circ \kappa_S. \quad (5.1)$$

We first show the existence of an R -bilinear form $\kappa_B^\theta \in \mathcal{L}_R^2(B)$ satisfying $(\kappa_B^\theta)_S = \theta^*(\kappa_S)$. The notation κ_B^θ indicates that, *a priori*, the form depends on θ (which in turn involves an S). According to (2.8) and Remark 2.10 the cocycle corresponding to the S -algebra isomorphism θ^{-1} is $u = \theta_2 \theta_1^{-1}$ and we have

$$\theta(B \otimes 1) = \{x \in \mathfrak{a} \otimes_k S : u((\text{Id}_{\mathfrak{a}} \otimes \alpha_1)(x)) = (\text{Id}_{\mathfrak{a}} \otimes \alpha_2)(x)\}. \quad (5.2)$$

We identify $B \subset B \otimes_R S$ via $b \mapsto b \otimes 1$ and claim that the restriction of $\theta^*(\kappa_S)$ to $B \times B$, which *a priori* takes values in S , actually takes values in R . In other words, for $b, b' \in B$ and $x = \theta(b), x' = \theta(b') \in \mathfrak{a} \otimes_k S$ we claim $\theta^*(\kappa_S)(b, b') = \kappa_S(x, x') \in R$. With $\alpha_i: S \rightarrow S''$ as before we have, using (5.1), (5.2) and the automorphism-invariance of $\kappa_{S''}$,

$$\begin{aligned} \alpha_1(\theta^*(\kappa_S)(b, b')) &= \alpha_1(\kappa_S(x, x')) = \kappa_{S''}((\text{Id}_{\mathfrak{a}} \otimes \alpha_1)(x), (\text{Id}_{\mathfrak{a}} \otimes \alpha_1)(x')) \\ &= \kappa_{S''}(u((\text{Id}_{\mathfrak{a}} \otimes \alpha_1)(x)), u((\text{Id}_{\mathfrak{a}} \otimes \alpha_1)(x'))) \\ &= \kappa_{S''}((\text{Id}_{\mathfrak{a}} \otimes \alpha_2)(x), (\text{Id}_{\mathfrak{a}} \otimes \alpha_2)(x')) \\ &= \alpha_2(\kappa_S(x, x')) = \alpha_2(\theta^*(\kappa_S)(b, b')). \end{aligned} \quad (5.3)$$

This shows that $\theta^*(\kappa_S)(b, b')$ belongs to the equalizer of α_1 and α_2 in S , but these are precisely the elements of R (viewed as elements of S). Hence the restriction κ_B^θ of $\theta^*(\kappa_S)$ to B is an R -bilinear form on B . Clearly, by its very definition, $(\kappa_B^\theta)_S = \theta^*(\kappa_S)$.

We next aim to show that κ_B^θ is independent of the trivialization θ . Thus, let S'/R be another faithfully flat extension, say with structure map $\alpha': R \rightarrow S'$, and let $\theta': B \otimes_R S' \rightarrow \mathfrak{a} \otimes_k S'$ be an S' -algebra isomorphism. By what we just have shown, we have an R -bilinear form $\kappa_B^{\theta'}$ satisfying $(\kappa_B^{\theta'})_{S'} = (\theta')^*(\kappa_S)$, and we claim $\kappa_B^\theta = \kappa_B^{\theta'}$. The vehicle to show this will be the algebra $T = S \otimes_R S'$ which we view in an obvious way as an S - and S' -algebra, say with structure maps β and β'

respectively. For simplicity we denote by $\rho: k \rightarrow R$ the structure map $\sigma_{R,k}$ of R . We have the following commutative diagram:

$$\begin{array}{ccccc}
 & & & S & \\
 & & \alpha & \nearrow & \beta \\
 k & \xrightarrow{\rho} & R & & T \\
 & & \alpha' & \searrow & \beta' \\
 & & & S' &
 \end{array}$$

We remind the reader that T is faithful flat over S , S' and R . Let $\xi: (\mathfrak{a} \otimes_{\alpha \circ \rho} S) \otimes_{\beta} T \rightarrow \mathfrak{a} \otimes_{\beta \circ \alpha \circ \rho} T = \mathfrak{a} \otimes_k T$ be the canonical T -linear algebra isomorphism, and define $\tilde{\theta}: \mathfrak{a} \otimes_{\beta \circ \alpha \circ \rho} T \rightarrow B \otimes_{\beta \circ \alpha} T$ by the composition of maps

$$\tilde{\theta}: \mathfrak{a} \otimes_{\beta \circ \alpha \circ \rho} T \xrightarrow{\xi^{-1}} (\mathfrak{a} \otimes_{\alpha \circ \rho} S) \otimes_{\beta} T \xrightarrow{(\theta \otimes \text{Id})^{-1}} (B \otimes_{\alpha} S) \otimes_{\beta} T \xrightarrow{\zeta} B \otimes_{\beta \circ \alpha} T.$$

Then $\tilde{\theta}^*$ maps $(\kappa_B^{\theta})_{\beta \circ \alpha} \in \mathcal{L}_T^2(B \otimes_{\beta \circ \alpha} T)$ onto a bilinear form in $\mathcal{L}_T^2(\mathfrak{a} \otimes_k T)$. In fact,

$$\begin{aligned}
 \tilde{\theta}^*((\kappa_B^{\theta})_{\beta \circ \alpha}) &= (\zeta \circ (\theta^{-1} \otimes \text{Id}) \circ \xi^{-1})^*((\kappa_B^{\theta})_{\beta \circ \alpha}) \\
 &= (\xi^{-1*} \circ (\theta^{-1} \otimes \text{Id})^* \circ \zeta^*)(\kappa_B^{\theta})_{\beta \circ \alpha} \\
 &\stackrel{(4.2)}{=} (\xi^{-1*} \circ (\theta^{-1} \otimes \text{Id})^*)((\kappa_B^{\theta})_{\alpha})_{\beta} \\
 &= (\xi^{-1*} \circ (\theta^{-1} \otimes \text{Id})^*)((\theta^* \kappa_{\alpha \circ \rho})_{\beta}) \\
 &\stackrel{(4.3)}{=} (\xi^{-1*} \circ (\theta^{-1} \otimes \text{Id})^* \circ (\theta \otimes \text{Id})^*)((\kappa_{\alpha \circ \rho})_{\beta}) \\
 &= \xi^{-1*}((\kappa_{\alpha \circ \rho})_{\beta}) \stackrel{(4.2)}{=} \kappa_{\beta \circ \alpha \circ \rho} = \kappa_T.
 \end{aligned}$$

We define $\tilde{\theta}': \mathfrak{a} \otimes_k T \rightarrow B_{\beta' \circ \alpha'} T$ by replacing $\theta \otimes \text{Id}$ by $\theta' \otimes \text{Id}$ in the definition of $\tilde{\theta}$ above. By symmetry $(\tilde{\theta}')^*(\kappa_B^{\theta'})_{\beta' \circ \alpha'} = \kappa_T$. Since $\tilde{\theta}^{-1} \circ \tilde{\theta}' \in \mathbf{Aut}(\mathfrak{a})(T)$, we get

$$(\kappa_B^{\theta'})_{\beta' \circ \alpha'} = (\tilde{\theta}')^{-1*}(\kappa_T) = \tilde{\theta}^{-1*}(\kappa_T) = (\kappa_B^{\theta})_{\beta \circ \alpha}$$

so that Lemma 4.2(c) implies $\kappa_B^{\theta} = \kappa_B^{\theta'}$. We are now justified to define $\kappa_B = \kappa_B^{\theta}$.

Finally, we use Lemma 5.3 to establish that κ_B is $\mathbf{Aut}(B)$ -invariant: κ is $\mathbf{Aut}(\mathfrak{a})$ -invariant $\Rightarrow \kappa_S$ is $\mathbf{Aut}(\mathfrak{a}_S)$ -invariant $\Rightarrow \theta^*(\kappa_S) = (\kappa_B)_S$ is $\mathbf{Aut}(B \otimes_{\alpha} S)$ -invariant $\Rightarrow \kappa_B$ is $\mathbf{Aut}(B)$ -invariant.

(b) κ_B is invariant since invariance of bilinear forms is stable under base ring extensions and, by Lemma 4.2(c), also under faithfully flat descent.

(c) The argument is analogous to that of (b), using Lemma 4.2(d) and the fact that being finitely presented is a property which is invariant under arbitrary base change and faithfully flat descent. \square

Corollary 5.5. *Let B be an S/R -form of $A = \mathfrak{a} \otimes_k R$ as in the descent setting (2.9). We further assume that \mathfrak{a} is finitely presented as a k -module and that $\kappa \in \text{IBF}_k(\mathfrak{a})$ is*

Aut(a)-invariant. Let κ_B be the R -bilinear form on B associated to κ as described in Theorem 5.4. If the IBF-principle holds for (\mathbf{a}, κ) , then the IBF-principle holds for (B, κ_B) as well.

Proof. By faithfully flat descent, the associated map $\overline{\kappa_B} : \mathbf{IBF}_R(B) \rightarrow R$ is an isomorphism as soon as the extended map $\overline{(\kappa_B)_S}$ has this property. By Lemma 4.6(a) applied to $\beta = \kappa_B$ we have $\overline{(\kappa_B)_S} = \overline{(\kappa_B)_S} \circ \nu$. Since ν is an isomorphism, we are reduced to showing that $\overline{(\kappa_B)_S}$ is an isomorphism. By the theorem, $(\kappa_B)_S = \theta^*(\kappa_S)$. From the definitions, the reader easily checks that $\overline{\theta^*(\kappa_S)} = \overline{\kappa_S} \circ \mathbf{IBF}(\theta)$. By functoriality $\mathbf{IBF}(\theta)$ is an isomorphism while $\overline{\kappa_S}$ is an isomorphism by Lemma 4.6(b). Hence $\overline{(\kappa_B)_S} = \overline{\kappa_S} \circ \mathbf{IBF}(\theta)$ is an isomorphism. This completes the proof. \square

6. Applications to Lie and Other Classes of Algebras

We now look in detail at our general results in some important special cases. Unless stated otherwise, we use our basic setting: k is a commutative associative unital ring and $R \in k\text{-alg}$. If f is an endomorphism of a finitely generated and projective R -module, we denote by $\text{tr}(f)$ its trace. For details on this notion, see, for example, [8, II, Sec. 4.3] or [13].

6.1. Lie algebras

We start by discussing the Killing form of a Lie algebra L , defined by $\kappa(l_1, l_2) = \text{tr}((\text{ad } l_1) \circ (\text{ad } l_2))$ for $l_i \in L$.

Proposition 6.1. *Let L be a Lie algebra over R whose underlying R -module is finitely generated and projective.*

(a) *The Killing form κ of L is an invariant and $\mathbf{Aut}(L)$ -invariant R -bilinear form. For any $S \in R\text{-alg}$ the base change κ_S is the Killing form of the Lie algebra $L \otimes_R S$.*

(b) *Suppose f is an α -semilinear automorphism of L for some $\alpha \in \text{Aut}_k(R)$. Then $\kappa(f(l_1), f(l_2)) = \alpha(\kappa(l_1, l_2))$ holds for $l_i \in L$.*

Proof. (a) The invariance of κ follows from $\text{tr}(fg) = \text{tr}(gf)$ for endomorphisms f, g of L . This identity also implies that κ is $\text{Aut}_R(L)$ -invariant: For $f \in \text{Aut}_R(L)$ we have $\text{ad}(f(l)) = f(\text{ad } l)f^{-1}$ and hence $\kappa(f(l_1), f(l_2)) = \text{tr}(f(\text{ad } l_1)(\text{ad } l_2)f^{-1}) = \kappa(l_1, l_2)$. The adjoint maps of the Lie S -algebra $L \otimes_R S$ are obtained by base change from the adjoint maps of L . Since the trace is invariant under base change, the Killing form of the Lie algebra $L \otimes_R S$ is the base change of κ by S . It then follows that κ is $\mathbf{Aut}(L)$ -invariant.

(b) We first prove an auxiliary formula. Namely, assume that g is an α^{-1} -semilinear endomorphism of L . Then

$$\text{tr}(fg) = \alpha(\text{tr}(gf)) \tag{6.1}$$

(observe that both fg and gf are R -linear). By descent it is sufficient to show (6.1) in case L is free of finite rank. Let F and G be matrices representing f and g in some R -basis of L . We denote by $\alpha \cdot G$ the matrix obtained from G by applying α to all its entries. Then fg and gf are represented by $F(\alpha \cdot G)$ and $G(\alpha^{-1} \cdot F)$ respectively, whence $\text{tr}(fg) = \text{tr}(F(\alpha \cdot G)) = \text{tr}((\alpha \cdot G)F) = \text{tr}(\alpha \cdot (G(\alpha^{-1} \cdot F))) = \alpha(\text{tr}(gf))$. We can now establish (b): $\kappa(f(l_1), f(l_2)) = \text{tr}(f(\text{ad } l_1 \text{ ad } l_2 f^{-1}))$. Applying (6.1) with $g = \text{ad } l_1 \text{ ad } l_2 f^{-1}$ shows that $\text{tr}(f(\text{ad } l_1 \text{ ad } l_2 f^{-1})) = \alpha(\text{tr}(\text{ad } l_1 \text{ ad } l_2 f^{-1} f)) = \alpha(\kappa(l_1, l_2))$. □

Corollary 6.2. *In the descent setting (2.9) suppose that \mathfrak{a} is a Lie algebra whose underlying k -module is finitely generated and projective. Let κ be the Killing form of \mathfrak{a} . Then*

- (a) *B is a finitely generated projective R -module and the unique R -bilinear form κ_B on B associated to κ in Theorem 5.4 is the Killing form of the Lie algebra B . If B is realized as an R -subalgebra of $\mathfrak{a} \otimes_k S$ as explained in (2.8),^f the form κ_B is the restriction of the Killing form κ_S of $\mathfrak{a} \otimes_k S$ to B .*
- (b) *If κ is nonsingular, then the Killing form of B is nonsingular.*
- (c) *If κ is nonsingular and \mathfrak{a} is a central k -algebra, then B is a central R -algebra, and $\text{IBF}_R(B)$ is free R -module of rank 1 admitting κ_B as a basis. In particular $\text{IBF}_R(B) = R\kappa_B$.*

Proof. (a) By Proposition 6.1, κ_S is the Killing form of the S -Lie algebra $\mathfrak{a} \otimes_k S$. The property of being finitely generated and projective is stable under arbitrary base change and faithfully flat descent. Hence the R -module B is finitely generated and projective. By Proposition 6.1, β_S is the Killing form of B_S . Let $\theta: B \otimes_R S \rightarrow \mathfrak{a} \otimes_k S$ be a trivialization. Since the isomorphism θ preserves Killing forms, we get $\beta_S = \theta^*(\kappa_S)$. Now (a) follows from the uniqueness assertion in Theorem 5.4.

(b) This follows from (a) and Lemma 4.2(d) (we remind the reader that every finitely generated projective module is finitely presented).

(c) By [25, Lemma 3.1] the S -algebra $\mathfrak{a} \otimes_k S$ is central. The faithfully flat descent reasoning of [11, Lemma 4.6(3)] then shows that the R -algebra B is a central. The last claim now follows from Corollary 3.8. □

At this point a very natural question arises: What are interesting examples of Lie algebras for which the IBF-principle 3.6 holds with respect to the Killing form? To convince the reader that (over rings) one cannot expect easy answers, we will look at one of the simplest and innocent looking Lie algebras.

Example 6.3 ($\mathfrak{sl}_2(k)$). Let $\mathfrak{sl}_2(k)$ be the Lie algebra of all traceless 2×2 -matrices with entries in our ring k . Its underlying module is free of rank 3, with the following

^fWhich is always possible up to R -isomorphism — this is the content of (2.8).

matrices forming the standard basis:

$$e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

A straightforward calculation shows that $\text{ibf}_k(\mathfrak{sl}_2(k))$ is spanned by

$$h \otimes e, e \otimes h, f \otimes h, h \otimes f, 2e \otimes e, 2f \otimes f, h \otimes h - 2f \otimes e, h \otimes h - 2e \otimes f.$$

Hence

$$\mathbf{IBF}_k(\mathfrak{sl}_2(k)) = ((k/2k)\overline{e \otimes e}) \oplus ((k/2k)\overline{f \otimes f}) \oplus \text{Span}_k\{\overline{e \otimes f}, \overline{f \otimes e}\}.$$

Consequently:

- If $2k = 0$, then $\mathbf{IBF}_k(\mathfrak{sl}_2(k))$ is free of rank 4, with basis $\{\overline{e \otimes e}, \overline{f \otimes f}, \overline{e \otimes f}, \overline{f \otimes e}\}$.
- If $2 \in k^\times$, i.e. $2 \in k$ is invertible, then $\mathbf{IBF}_k(\mathfrak{sl}_2(k))$ is free of rank 1, e.g., with basis $\{\overline{h \otimes h}\}$.

Using the isomorphism (3.6) and the description of $\mathbf{IBF}(\mathfrak{sl}_2(k))$ above, we can define an invariant bilinear form $\gamma \in \mathbf{IBF}_k(\mathfrak{sl}_2(k))$, sometimes called the *normalized Killing form* or the *normalized invariant form*, by

$$\gamma(e, e) = 0 = \gamma(f, f), \quad \gamma(e, f) = 1 = \gamma(f, e).$$

Note that $\gamma(h, h) = 2$ and that all other values of γ on the standard basis of $\mathfrak{sl}_2(k)$ are zero, in particular γ is symmetric. The description of $\mathbf{IBF}_k(\mathfrak{sl}_2(k))$ above implies that:

If 2 is not a zero divisor in k, then $\mathbf{IBF}_k(\mathfrak{sl}_2(k)) = k\gamma$ is free of rank 1.

Moreover, by calculating the discriminant of γ one obtains:

$$\gamma \text{ is nonsingular} \Leftrightarrow 2 \in k^\times \Leftrightarrow \text{the IBF-principle holds for } (\mathfrak{sl}_2(k), \gamma).$$

In this case, γ is $\mathbf{Aut}(\mathfrak{sl}_2(k))$ -invariant, which can be seen by noting that $(\text{ad } x)^3 - 2\gamma(x, x)\text{ad } x = 0$ is the generic minimal polynomial of $\mathfrak{sl}_2(k)$.

It is straightforward that 12γ is the Killing form of $\mathfrak{sl}_2(k)$. In particular, the Killing form vanishes if $2k = 0$ (not surprising since then $\mathfrak{sl}_2(k)$ is a 2-step nilpotent Lie algebra) or if $3k = 0$ (somewhat surprising since $\mathfrak{sl}_2(k)$ is a simple Lie algebra when k is a field of characteristic 3). The conclusion is that for the setting of this paper the normalized Killing form γ is better behaved than the Killing form itself. A case in point is a revised version of Corollary 6.2 for $\mathfrak{a} = \mathfrak{sl}_2(k)$ and $2 \in k^\times$. Since $\mathfrak{sl}_2(k)$ is then central,[§] the proof of Corollary 6.2 shows that:

- If $2 \in k^\times$, any S/R -form B of $\mathfrak{sl}_2(R)$ is central and has a nonsingular invariant bilinear form β (not necessarily the Killing form), for which $\mathbf{IBF}_R(B) = R\beta$ is free of rank 1.

[§]Centrality only requires that 2 not be a zero divisor.

It is instructive to summarize what we have shown above for the special case $k = \mathbb{Z}$.

- The k -module $\mathbf{IBF}_{\mathbb{Z}}(\mathfrak{sl}_2(\mathbb{Z}))$ is neither projective nor cyclic, in particular, the IBF-principle does not hold for $\mathfrak{sl}_2(\mathbb{Z})$. Yet $\mathbf{IBF}_{\mathbb{Z}}(\mathfrak{sl}_2(\mathbb{Z})) = \mathbb{Z}\gamma$ is free of rank 1.
- All invariant bilinear forms of $\mathfrak{sl}_2(\mathbb{Z})$ are symmetric (even though $\mathfrak{sl}_2(\mathbb{Z})$ is not perfect, cf. Remark 3.4).
- All nonzero invariant bilinear forms of $\mathfrak{sl}_2(\mathbb{Z})$ are nondegenerate, but none of them is nonsingular.

Remark 6.4 (Generalizations of Example 6.3). We note that Example 6.3 can be generalized by replacing $\mathfrak{sl}_2(k) = \mathfrak{sl}_2(\mathbb{Z}) \otimes_{\mathbb{Z}} k$ by $\mathcal{G} \otimes_{\mathbb{Z}} k$ where \mathcal{G} is the Lie algebra of a split simple simply connected Chevalley–Demazure group scheme. In terms of Lie algebras, \mathcal{G} is a Chevalley order of a split simple Lie algebra $(\mathfrak{g}, \mathfrak{h})$ over \mathbb{Q} , say with root system Δ , which is compatible with the root space decomposition of $(\mathfrak{g}, \mathfrak{h})$ and satisfies $\mathcal{G} \cap \mathfrak{h} = \text{Span}_{\mathbb{Z}}\{h_{\alpha} : \alpha \in \Delta\}$ (with the standard notation). In this setting the existence of an invariant bilinear form γ as above follows from [29, 12]. It is uniquely determined by the condition $\gamma(h_{\alpha}, h_{\alpha}) = 2$ for any long root α . Details will be left to the reader.

In what follows we restrict our presentation to base fields of characteristic 0. To abide by standard notation we denote our algebra \mathfrak{a} , which is now a finite-dimensional semisimple Lie algebra defined over a field k of characteristic 0, by \mathfrak{g} . We are interested in twisted forms of $\mathfrak{g} \otimes_k R$ for some $R \in k\text{-alg}$. In the case when k is algebraically closed, \mathfrak{g} is simple and R is the Laurent polynomial ring $k[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$, the twisted forms in question are related to the affine Kac–Moody Lie algebras (the case $n = 1$) and more generally to multiloop algebras (see [2, 11, 22, 25] and Sec. 7 for further details and references).

Theorem 6.5. *Let \mathfrak{g} be a finite-dimensional semisimple Lie algebra over a field k of characteristic 0. Let B be a twisted form of $\mathfrak{g} \otimes_k R$, split by a faithfully flat extension S/R .*

(a) *Then B is a finitely generated projective R -module and perfect as a Lie algebra. The Killing form of the R -algebra B coincides with the bilinear form κ_B associated to the Killing form κ of \mathfrak{g} in Theorem 5.4. In particular the Killing form of B is nonsingular and $\mathbf{Aut}(B)$ -invariant. If B is realized as an R -subalgebra of $\mathfrak{g} \otimes_k S$, the Killing form of B is the restriction of the Killing form $\kappa_{\mathfrak{g} \otimes_k S}$ to B .*

(b) *Assume henceforth that \mathfrak{g} is central, hence central-simple. Then B is a central R -algebra, $\mathbf{IBF}_R(B)$ is a free R -module of rank 1 admitting κ_B as a basis, and (B, κ_B) satisfies the IBF-principle 3.6.*

Proof. With the exception of the perfectness statement, (a) and the first part of (b) is a re-statement of Corollary 6.2. But being perfect is a property (of arbitrary algebras) which is stable under arbitrary base change and faithfully flat descent. Since $B \otimes_R S \simeq \mathfrak{g} \otimes_k S$ and the latter is perfect, B is perfect. That (B, κ_B) satisfies

the IBF-principle is a special case of Corollary 5.5 keeping in mind that (\mathfrak{g}, κ) satisfies the IBF-principle in view of Proposition 3.9. \square

Remark 6.6. In the special case when $R = S$ and $B = \mathfrak{g} \otimes_k R$, our result says that

$$\mathbf{IBF}(\mathfrak{g} \otimes_k R) \simeq R \simeq \mathbf{IBF}_k(\mathfrak{g}) \otimes_k R. \quad (6.2)$$

This formula is also a special case of [33, Theorem 4.1] which, using methods from Lie algebra homology, determines the “predual” of the space of symmetric bilinear forms of a Lie algebra of type $L \otimes_k R$ for k a field of characteristic $\neq 2$ and L any Lie algebra over k . Assuming for comparison reasons that L is perfect, we know (Remark 3.4) that all invariant bilinear maps are symmetric and thus [33, Theorem 4.1] becomes (6.2) for Lie algebras of type $L \otimes_k R$. We emphasize that the approach of [33] *cannot* be applied to the case of twisted forms of $\mathfrak{g} \otimes_k R$. Developing method that would apply to these algebras was the original motivation for our work. As already observed, such twisted algebras already arise in the affine Kac–Moody setting and are crucial for EALA theory.

An immediate consequence of (6.2) is that every invariant bilinear form $\beta \in \mathbf{IBF}_{(R,k)}(\mathfrak{g} \otimes_k R) = \mathbf{IBF}_k(\mathfrak{g} \otimes_k R)$ has the form $\varphi \circ \kappa_R$ for a unique $\varphi \in R^* = \text{Hom}_k(R, k)$, i.e. $\beta(x_1 \otimes r_1, x_2 \otimes r_2) = \kappa(x_1, x_2) \varphi(r_1 r_2)$ for $x_i \in \mathfrak{g}$ and $r_i \in R$. This latter fact has recently been re-proven in [18, Lemma 2.3] in case k is an algebraically closed field of characteristic 0, using the structure theory of \mathfrak{g} .

The untwisted case was, out of necessity, the first objective of our work. The methods to be developed, however, had to be compatible with descent theory so that results about twisted algebras could be obtained. In retrospect, we “knew” that the functor on k -spaces $\mathbf{IBF}_k(\mathfrak{g} \otimes_k S, -)$ is represented by S (one can reinterpret [33] or [18] this way). But how does one recover S from $\mathfrak{g} \otimes_k S$? The answer is as its centroid. By descent, the centroid of B is in this case naturally isomorphic to R . Our result shows that the representability of $\mathbf{IBF}_k(B, -)$ in terms of the centroid is indeed the correct point of view.

Remark 6.7. For crucial use in [27], we note the following. Since the IBF-principle holds by Theorem 6.5(b), composing the isomorphism (3.9) with the inverse of the isomorphism given in Lemma 3.7, we have an isomorphism $\text{Hom}_k(R, V) \rightarrow \text{Ctd}_R(B, \text{Hom}_k(B, V))$, $\varphi \mapsto \tilde{\varphi}$ such that $\tilde{\varphi}(b)(b') = \varphi(\kappa_B(b, b'))$.

6.2. Unital algebras

In this subsection we will discuss invariant bilinear forms of unital algebras B defined over some $R \in k\text{-alg}$. To do so, we will use the *associator module* and *commutator module* defined for an arbitrary algebra B by

$$(B, B, B) = \text{Span}_{\mathbb{Z}}\{(a, b, c) : a, b, c \in B\} \quad \text{and} \quad [B, B] = \text{Span}_{\mathbb{Z}}\{[a, b] : a, b \in B\}$$

respectively, where $(a, b, c) = (ab)c - a(bc)$ is the associator and $[a, b] = ab - ba$ is the commutator in B .^h It is immediate that (B, B, B) and $[B, B]$ are R -submodules of B . We define

$$\mathfrak{ac}(B) = (B, B, B) + [B, B] \quad \text{and} \quad \mathbb{A}\mathbb{C}(B) = B/\mathfrak{ac}(B).$$

Let $1_B \in B$ be the identity element of B . Thus $b1_B = b = 1_B b$ for all $b \in B$. A unital algebra is perfect, whence $\mathbf{IBF}_{(R,k)}(B; V) = \mathbf{IBF}_k(B; V)$ for any k -module V by Remark 3.4.

For convenience for the remainder of this section we will denote the identity element of B by 1.

Lemma 6.8. *Let B be a unital R -algebra. Then the multiplication map $\mu: B \otimes_R B \rightarrow B$, $\mu(a \otimes b) = ab$, induces an isomorphism*

$$\bar{\mu}: \mathbf{IBF}_R(B) \rightarrow \mathbb{A}\mathbb{C}(B), \quad \bar{\mu}(\overline{a \otimes b}) = \overline{ab}$$

with inverse given by $\bar{a} \mapsto \overline{1 \otimes a} = \overline{a \otimes 1}$. Hence, for any k -module V the natural map

$$\mathrm{Hom}_k(\mathbb{A}\mathbb{C}(B), V) \rightarrow \mathbf{IBF}_k(B; V),$$

which assigns to $\varphi \in \mathrm{Hom}_k(\mathbb{A}\mathbb{C}(B), V)$ the bilinear function $(a, b) \mapsto \varphi(\overline{ab})$, is an isomorphism of R -modules. Its inverse is given by assigning to β the linear function $\bar{b} \mapsto \beta(b, 1)$, where $b \in B$.

Proof. By (3.3), $\mathbf{ibf}_R(B)$ is spanned by elements of the form $ab \otimes c - a \otimes bc$ and $a \otimes b - b \otimes a$. It is clear that $\mu(\mathbf{ibf}_R(B)) = \mathfrak{ac}(B)$. Hence $\bar{\mu}$ is well-defined and surjective. Let $\nu: B \rightarrow B \otimes_k B$ be defined by $\nu(a) = 1 \otimes a$. Then we have

$$\nu((a, b, c)) = 1 \otimes (ab)c - 1 \otimes a(bc) \equiv ab \otimes c - a \otimes bc \equiv 0 \pmod{\mathbf{ibf}_R(B)}, \quad \text{and}$$

$$\nu([a, b]) = 1 \otimes ab - 1 \otimes ba \equiv a \otimes b - b \otimes a \equiv 0 \pmod{\mathbf{ibf}_R(B)}.$$

We thus get a well-defined k -linear map $\bar{\nu}: \mathbb{A}\mathbb{C}(B) \rightarrow \mathbf{IBF}_R(B)$ satisfying $\bar{\nu}(\bar{b}) = \overline{b \otimes 1} = \overline{1 \otimes b}$. Because of $(\nu \circ \mu)(a \otimes b) = 1 \otimes ab = (1 \otimes ab - 1a \otimes b) + a \otimes b \equiv a \otimes b \pmod{\mathbf{ibf}_R(B)}$, we have $\bar{\nu} \circ \bar{\mu} = \mathrm{Id}_{\mathbf{IBF}(B)}$, proving injectivity and thus bijectivity of $\bar{\mu}$. Under the isomorphism $\bar{\mu}$, the universal bilinear map $\beta_{\mathrm{uni}}: B \times B \rightarrow \mathbf{IBF}_R(B)$ becomes $\beta_{\mathrm{uni},u}: B \times B \rightarrow \mathbb{A}\mathbb{C}(B)$ with $\beta_{\mathrm{uni},u}(a, b) = \overline{ab}$. In view of (3.7) this implies the last claim. \square

^hIf B happens to be a Lie algebra, the commutator as defined here is twice the Lie algebra product. This notational conflict should not cause any problems since in the following we will employ the notation $[a, b]$ for non-Lie algebras only.

Corollary 6.9. *Let B be a unital R -algebra and assume that $B = Rb_0 \oplus \mathfrak{ac}(B)$ for some $b_0 \in B$ where Rb_0 is free with basis $\{b_0\}$. Let $\pi: B \rightarrow R$ be defined by $b = \pi(b)b_0 \oplus b_{\mathfrak{ac}}$ where $b_{\mathfrak{ac}} \in \mathfrak{ac}(B)$, and define*

$$\beta_0: B \times B \rightarrow R, \quad \beta_0(a, b) = \pi(ab).$$

Then $\beta_0 \in \text{IBF}_R(B)$, and (B, β_0) satisfies the IBF-principle. Furthermore

$$\tilde{\beta}_0: \mathbb{A}\mathbb{C}(B) \rightarrow R, \quad \bar{b} \mapsto \beta_0(b, 1)$$

is a well-defined R -module isomorphism.

Proof. The proof is straightforward. □

6.3. Azumaya algebras

This subsection fits within the descent setting (2.9): k is base ring, $R \in k\text{-alg}$ is flat as a k -module (for example $k = R$), and $S \in R\text{-alg}$ is a faithfully flat R -module. We let $\mathfrak{a} = M_n(k)$. Then our S/R -form B of $M_n(k) \otimes_k R = M_n(R)$ is an Azumaya algebra over R of constant rank n^2 .

We start by recording some facts about \mathfrak{a} . As a unital algebra, \mathfrak{a} is perfect. It is also well-known that \mathfrak{a} is central. Moreover, \mathfrak{a} has a natural invariant bilinear form, the trace form κ defined by

$$\kappa(x, y) = \text{tr}(xy),$$

where this last is the usual trace of the matrix xy . It is easy to see (using the standard dual basis of the elementary matrices E_{ij}) that κ is nonsingular. Moreover, κ is $\mathbf{Aut}(\mathfrak{a})$ -invariant. Indeed, since $\mathfrak{a} \otimes_k K = M_n(K)$ for any $K \in k\text{-alg}$, it suffices to verify that κ is automorphism-invariant. Thus let $\sigma \in \text{Aut}_k(\mathfrak{a})$ and $x, y \in \mathfrak{a}$. To show that xy and $\sigma(x)\sigma(y)$ have the same trace, it is enough to prove that for all $\mathfrak{p} \in \text{Spec}(k)$ the two elements $(xy)_{\mathfrak{p}}$ and $(\sigma(x)\sigma(y))_{\mathfrak{p}}$ of $M_n(k_{\mathfrak{p}})$ have the same trace. Clearly $(xy)_{\mathfrak{p}} = x_{\mathfrak{p}}y_{\mathfrak{p}}$ and $(\sigma(x)\sigma(y))_{\mathfrak{p}} = \sigma_{\mathfrak{p}}(x_{\mathfrak{p}})\sigma_{\mathfrak{p}}(y_{\mathfrak{p}})$ where $\sigma_{\mathfrak{p}} = \sigma \otimes \text{Id}_{k_{\mathfrak{p}}}$. We may therefore assume that k is a local ring. But then, by the Skolem–Noether theorem for local rings [13, IV, Corollary 1.3], σ is given by conjugation by an invertible matrix $M \in \text{GL}_n(k)$, whence $\sigma(x)\sigma(y) = MxyM^{-1}$, and so clearly xy and $\sigma(x)\sigma(y)$ have the same trace. We also have

$$\mathfrak{a} = kE_{11} \oplus [\mathfrak{a}, \mathfrak{a}], \quad [\mathfrak{a}, \mathfrak{a}] = \{x \in \mathfrak{a} : \text{tr}(x) = 0\} = \mathfrak{ac}(\mathfrak{a})$$

since any $x = \sum_{i,j} x_{ij}E_{ij}$ can be uniquely written as

$$x = \left(x_{11} + \sum_{1 < i} x_{ii} \right) E_{11} + \sum_{1 < i} x_{ii}(E_{ii} - E_{11}) + \sum_{i \neq j} x_{ij}E_{ij} \quad (6.3)$$

and $[\mathfrak{a}, \mathfrak{a}]$ is spanned by matrices of type $[E_{ii}, E_{ij}] = E_{ij}$ and $[E_{ij}, E_{ji}] = E_{ii} - E_{jj} = (E_{ii} - E_{11}) - (E_{jj} - E_{11})$ for $i \neq j$. Formula (6.3) implies that the trace form κ is the bilinear form of Corollary 6.9. Hence (\mathfrak{a}, κ) satisfies the IBF-principle.

Theorem 6.10. *Let B be an S/R -form of $M_n(R)$ and let κ_B be the bilinear form associated to the trace form κ of $M_n(k)$ in Theorem 5.4.*

- (a) *Then κ_B is a nonsingular, invariant and $\mathbf{Aut}(B)$ -invariant bilinear form and a basis of $\mathbf{IBF}_R(B)$.*
- (b) *$\mathbf{IBF}_R(B) \simeq \mathbb{A}\mathbb{C}(B) \simeq R$, and the map $\mathbf{IBF}_R(B) \mapsto R, \overline{b \otimes b'} \mapsto \kappa_B(b, b')$ is an isomorphism of R -modules. Hence (B, κ_B) satisfies the IBF-principle 3.6.*
- (c) *If B is realized as an R -subalgebra of $M_n(S)$, see Remark 2.10, κ_B coincides with the restriction of the trace form of $M_n(S)$ to B .*

Proof. Parts (a) and (b) follow from Theorem 5.4, Corollary 5.5 and Corollary 6.9. For the proof of (c) one uses the automorphism invariance of the trace of $M_n(S'')$ and the reasoning in (5.3) to conclude that the restriction λ of the trace form of $M_n(S)$ to B has values in R . Since λ_S is the trace form of $M_n(S)$ and thus coincides with $(\kappa_B)_S$, we get $\lambda = \kappa_B$ from uniqueness in Theorem 5.4 (or from Lemma 4.2(c)). \square

Remark 6.11. The form κ_B is, by definition, nothing but the reduced trace form of the Azumaya algebra B as defined in [13]. This proves (without the construction of the characteristic polynomial as done in [13]) that the reduced trace form, which *a priori* takes values in S , does take values in R .

Corollary 6.12. *Every Azumaya algebra B over R has a nonsingular, invariant and $\mathbf{Aut}(B)$ -invariant bilinear form κ_B such that (B, κ_B) satisfies the IBF-Principle 3.6. In particular, $\mathbf{IBF}_R(B)$ is a free R -module with basis $\{\kappa_B\}$.*

Proof. If B has constant rank, then B is an S/R -form of some $M_n(R)$ and the result follows from Theorem 6.10. In general, we can decompose the identity element 1_R of R into a sum $1_R = e_1 + \cdots + e_s$ of orthogonal idempotents $e_i \in R$ such that $B = B_1 \boxplus \cdots \boxplus B_s$ is a direct product of ideals $B_i = e_i B$, each B_i is an Azumaya algebra of constant rank ρ_i over $R_i = e_i R$, and $\rho_i \neq \rho_j$ for $i \neq j$ (if $\rho_i = \rho_j$ then we replace e_i, e_j by $e_i + e_j$). We then define κ_B as the orthogonal sum of the forms κ_{B_i} constructed previously. Nonsingularity follows from $\text{Hom}_R(B_1 \boxplus \cdots \boxplus B_s, R) \simeq \bigoplus_{i=1}^s \text{Hom}_{R_i}(B_i, R_i)$ and the nonsingularity of the κ_{B_i} . Finally, $\mathbf{Aut}(B)$ -invariance holds since the decomposition $B = B_1 \boxplus \cdots \boxplus B_s$ is preserved under base ring extensions and automorphisms. \square

6.4. Octonion algebras

As in the previous subsection, k here is an arbitrary base ring and $R \in k\text{-alg}$. Following [7, 17, 24] we call an algebra B over R an *octonion algebra* if its underlying R -module is projective of constant rank 8, contains an identity element 1_B , and admits a quadratic form $n_B: B \rightarrow R$, the *norm* of B , satisfying the following two

conditions.

- (i) The associated bilinear form $n_B : B \times B \rightarrow R$, $n_B(a, b) = n_B(a + b) - n_B(a) - n_B(b)$, is nonsingular, and
- (ii) $n_B(ab) = n_B(a)n_B(b)$ holds for all $a, b \in B$.

For an octonion algebra B the linear form $t_B = n_B(1_B, -)$ is called the *trace of B* . An example of an octonion algebra is the algebra $\text{Zor}(R)$ of *Zorn vector matrices*, defined on the R -module

$$Z = \text{Zor}(R) = \begin{bmatrix} R & R^3 \\ R^3 & R \end{bmatrix}$$

with product

$$\begin{bmatrix} \alpha_1 & u \\ x & \alpha_2 \end{bmatrix} \begin{bmatrix} \beta_1 & v \\ y & \beta_2 \end{bmatrix} = \begin{bmatrix} \alpha_1\beta_1 - {}^tuy & \alpha_1v + \beta_2u + x \times y \\ \beta_1x + \alpha_2y + u \times v & -{}^txv + \alpha_2\beta_2 \end{bmatrix}$$

for $\alpha_i, \beta_i \in R$ and $u, v, x, y \in R^3$. Here tuy and $x \times y$ are the usual scalar and vector product of vectors in R^3 . For this octonion algebra and $a = \begin{bmatrix} \alpha_1 & u \\ x & \alpha_2 \end{bmatrix} \in Z$ one has

$$1_Z = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad n_Z(a) = \alpha_1\alpha_2 + {}^tux, \quad \text{tr}_Z(a) = \alpha_1 + \alpha_2.$$

For our approach to octonion algebras it is important that an R -algebra B is an octonion algebra if and only if there exists a faithfully flat (even faithfully flat and étale) $S \in R\text{-alg}$ such that $B \otimes_R S \simeq \text{Zor}(S)$ (see [17, Corollary 4.11]). The algebra $\text{Zor}(R)$ is referred to as *split octonions*.

Theorem 6.13. *Let B be an octonion algebra over R .*

- (a) B is a central R -algebra satisfying $\text{ac}(B) = (B, B, B) = [B, B]$.
- (b) The bilinear form $\tau : B \times B \rightarrow R$, defined by $\tau(x, y) = \text{tr}(xy)$, is an invariant, nonsingular and $\mathbf{Aut}(B)$ -invariant R -bilinear form. It coincides with the form τ_B associated in Theorem 5.4 to the bilinear form τ of $\text{Zor}(R)$ with respect to any splitting $B \otimes_R S \simeq \text{Zor}(S)$ of B .
- (c) (B, τ) satisfies the IBF-principle.

Proof. The algebra B fits into our descent setting with $k = R$ and $\mathfrak{a} = \text{Zor}(R)$. We first prove all assertions for \mathfrak{a} . Straightforward calculations (admittedly tedious in the case of the associator module) show that:

- (i) $[\mathfrak{a}, \mathfrak{a}] = (\mathfrak{a}, \mathfrak{a}, \mathfrak{a}) = \text{ac}(\mathfrak{a})$, $\mathfrak{a} = \text{Re} \oplus \text{ac}(\mathfrak{a})$ for $e = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, \mathfrak{a} is central.
- (ii) For a, b as in the product formula above we have $\tau(a, b) = \alpha_1\beta_1 + \alpha_2\beta_2 - {}^tuy - {}^txv$. Hence τ is nonsingular and symmetric.
- (iii) $\tau(\text{ac}(\mathfrak{a})) = 0$, $\tau(e) = 1$, whence τ is the bilinear form associated to the decomposition $\mathfrak{a} = \text{Re} \oplus \text{ac}(\mathfrak{a})$ in Corollary 6.9.

This corollary now implies that (\mathfrak{a}, τ) satisfies the IBF-principle. To establish that τ is $\mathbf{Aut}(\mathfrak{a})$ -invariant, we recall that since \mathfrak{a} is a quadratic algebra the trace linear form tr of \mathfrak{a} is uniquely determined by the unital algebra \mathfrak{a} (see [24, Lemma 1.2]). For any extension $K \in k\text{-alg}$, the base change τ_K is therefore the bilinear form $\tau_{\mathfrak{a} \otimes_R K}$ of the K -algebra $\mathfrak{a} \otimes_R K = \text{Zor}(K)$. Uniqueness of the trace then implies that tr_K is $\text{Aut}_K(\mathfrak{a}_K)$ -invariant.

We now consider an arbitrary octonion algebra B over R and choose a faithfully flat extension $S \in R\text{-alg}$ such that $B \otimes_R S \simeq \text{Zor}(S)$ as S -algebras. In the first part of the proof we have established all claims for $\mathfrak{a} = \text{Zor}(R)$, whence also for $\text{Zor}(S)$. The assertions in (a) now hold for B since they are all preserved by faithfully flat descent. In (b) it suffices to establish the second part, but this follows from the fact that the base change of the trace form τ of B to S is the trace form of $\text{Zor}(S)$. Part (c) is a special case of Corollary 5.5. □

Remark 6.14 (Quadratic algebras). The experts will undoubtedly have noticed that the automorphism-invariance of the bilinear form κ_B comes from the fact that octonions are quadratic algebras, see, e.g., [24, 1.1]. Hence our techniques can also be applied to certain quadratic algebras whose trace forms are invariant.

6.5. *Alternative algebras*

We consider alternative algebras, always assumed to be unital, over some base ring R . Recall (see [7]) that an alternative algebra B is called *separable* if for every algebraically closed field K in $R\text{-alg}$ the K -algebra $B \otimes_R K$ is finite-dimensional and a direct sum of simple ideals. Equivalently, the unital universal multiplication envelope of B is a separable associative algebra. By [7, Proposition 2.11], an R -algebra B is central separable and alternative if and only if $R = R_1 \boxplus R_2$ is a direct sum of two ideals such that $B_1 = R_1 B$ is an Azumaya algebra over R_1 and $B_2 = R_2 B$ is an octonion algebra over R_2 . It is now straightforward to extend the results of Secs. 6.3 and 6.4 to central separable alternative algebras. We leave the details to the reader and only mention the following.

Corollary 6.15. *A central separable alternative B over R has a nonsingular invariant bilinear form κ_B such that $\text{IBF}_R(B)$ is a free R -module with basis $\{\kappa_B\}$.*

6.6. *Jordan algebras*

Central separable Jordan algebras over rings R containing $\frac{1}{2}$ (see [6, 15]) are another class of algebras to which our results apply.

By definition, a unital Jordan algebra J over R is separable if and only if $J \otimes_R K$ is finite-dimensional semisimple for all fields $K \in R\text{-alg}$. A central separable Jordan algebra J is generically algebraic (see [16, Example 2.4(d)]). Let $\text{tr} \in \text{Hom}_R(J, R)$ be its generic trace. By Proposition 2.7 of [16] the associated bilinear form τ , defined by $\tau(a, b) = \text{tr}(ab)$, is invariant, $\text{Aut}(J)$ -invariant and commutes with extensions and

faithfully flat descent. Since separability and being generically algebraic is invariant under base ring extensions, τ is in fact $\mathbf{Aut}(J)$ -invariant. By [14, Corollary 16.16], τ is nondegenerate for separable Jordan algebras over fields. From Lemma 6.16 we then get that τ is nonsingular.

Lemma 6.16. *Let M be a finitely generated projective R -module and let $\beta \in \mathcal{L}_R^2(M)$. Then β is nonsingular if and only if β_K is nondegenerate for all $K \in k\text{-alg}$ which are fields.*

Proof. This is an application of [9, II, Sec. 3.3, Théorème 1, Sec. 3.2, Corollaire de la Prop. 6 and Sec. 5.3, Théorème 2]. \square

Thus, in view of Corollary 3.8 we have the following.

Theorem 6.17. *The generic trace form τ of a central separable Jordan algebra J over a ring R containing $\frac{1}{2}$ is an invariant nonsingular and $\mathbf{Aut}(J)$ -invariant bilinear form, and $\mathbf{IBF}_R(J)$ is a free R -module admitting $\{\tau\}$ as a basis.*

We leave it to the interested reader to look into the following possible improvement of this last result.

Question 6.18. For J as in Proposition 6.17, is the R -module $\mathbf{IBF}_R(J)$ projective?

Since J is central and therefore a faithful R -module, we have $\kappa_J(J, J) = R$ by [10, I, Corollary 1.10]. Hence, if the question has a positive answer, Proposition 3.9 applies and yields that (J, τ) satisfies the IBF-principle. In particular, Theorem 6.17 then becomes a corollary.

7. Graded Invariant Bilinear Forms

In this section we classify graded invariant bilinear forms, which are particularly important for infinite-dimensional Lie theory. We will therefore concentrate on these (except for preliminaries considerations), and leave the extension to other classes of algebras to the reader.

We begin with some generalities about gradings and graded forms. Unless specified otherwise, we continue with our standard setting: k is a base ring and B is an arbitrary R -algebra for some $R \in k\text{-alg}$. Throughout, Λ is an abelian group.

Definition 7.1 (Graded algebras and graded invariant bilinear forms). A Λ -graded algebra is a pair (C, \mathcal{C}) consisting of a k -algebra C and a family $\mathcal{C} = (C^\lambda)_{\lambda \in \Lambda}$ of k -submodules C^λ of C satisfying $C = \bigoplus_{\lambda \in \Lambda} C^\lambda$ and $C^\lambda C^\mu \subset C^{\lambda+\mu}$ for all $\lambda, \mu \in \Lambda$. We will say that a k -algebra C is Λ -graded if (C, \mathcal{C}) is a Λ -graded algebra for some family \mathcal{C} . We point out that it is allowed that some of the homogeneous submodules C^λ vanish. If C is a unital algebra, then necessarily $1_C \in C^0$.

Assume C is Λ -graded. We call $\kappa \in \mathcal{L}_k^2(C)$ a *graded bilinear form* if $\kappa(C^\lambda, C^\mu) = 0$ whenever $\lambda + \mu \neq 0$.

Definition 7.2 (Graded S/R -forms of algebras). In the following we assume that $R \in k\text{-alg}$ is Λ -graded, say $R = \bigoplus_{\lambda \in \Lambda} R^\lambda$. An R -algebra B is then called a Λ -graded R -algebra if $B = \bigoplus_{\lambda \in \Lambda} B^\lambda$ is Λ -graded as a k -algebra and the Λ -gradings of R and B are compatible in the sense that $R^\lambda B^\mu \subset B^{\lambda+\mu}$ for all $\lambda, \mu \in \Lambda$. For example, for any k -algebra \mathfrak{a} the R -algebra $\mathfrak{a} \otimes_k R$ is canonically a Λ -graded R -algebra by defining the λ -homogeneous submodule $(\mathfrak{a} \otimes_k R)^\lambda = \mathfrak{a} \otimes_k R^\lambda$.

In the descent setting (2.9) we suppose that $R \in k\text{-alg}$ is Λ -graded and that $S \in R\text{-alg}$ is a Λ -graded R -algebra. We view $\mathfrak{a} \otimes_k S$ with its canonical Λ -grading. An S/R -form B of $\mathfrak{a} \otimes_k R$ is called *graded* if B is a Λ -graded R -algebra and there exists an S -algebra isomorphism $\theta: B \otimes_R S \rightarrow \mathfrak{a} \otimes_k S$ which respects the gradings: $\theta(b^\lambda \otimes s^\mu) \in \mathfrak{a} \otimes_k S^{\lambda+\mu}$ for $b^\lambda \in B^\lambda$ and $s^\mu \in S^\mu$.

We now specialize to Lie algebras and derive a graded version of Theorem 6.5. Following the notation used in Theorem 6.5 we change \mathfrak{a} to \mathfrak{g} .

Proposition 7.3. *Let \mathfrak{g} be a finite-dimensional semisimple Lie algebra over a field k of characteristic 0 with Killing form κ . Assume that $R \in k\text{-alg}$ and $S \in R\text{-alg}$ are Λ -graded, S/R is faithfully flat and B is a graded S/R -form of $\mathfrak{g} \otimes_k R$.*

- (a) *If κ_B is the form attached to κ in Theorem 5.4, then κ_B is the Killing form of the R -algebra B and satisfies $\kappa_B(B^\lambda, B^\mu) \subset R^{\lambda+\mu}$ for all $\lambda, \mu \in \Lambda$.*
- (b) *If \mathfrak{g} is central (and hence simple), every graded invariant bilinear form $\beta \in \text{IBF}_k(B)$ can be written in the form $\varphi \circ \kappa_B$ for a unique $\varphi \in \{\varphi \in R^* : \varphi(R^\lambda) = 0 \text{ for } \lambda \neq 0\} \simeq (R^0)^*$.*

Proof. (a) That κ_B is the Killing form of B , was established in Corollary 6.2. Since there exists an S -algebra isomorphism $\theta: B \otimes_R S \rightarrow \mathfrak{g} \otimes_k S$ respecting the gradings, there is no harm to assume that $B \subset \mathfrak{g} \otimes_k S$ with $B^\lambda = B^\lambda \otimes 1_S \subset \mathfrak{g} \otimes S^\lambda$. Recall from Corollary 6.2 that $\kappa_B = \kappa_{\mathfrak{g} \otimes_k S} | B \times B$ where $\kappa_{\mathfrak{g} \otimes_k S}$ is the Killing form of the S -algebra $\mathfrak{g} \otimes_k S$, and that $\kappa_{\mathfrak{g} \otimes_k S}$ coincides with the base change κ_S of κ by S . Because $\kappa_S(\mathfrak{g} \otimes_k S^\lambda, \mathfrak{g} \otimes_k S^\mu) \subset S^{\lambda+\mu}$, it suffices to show that $R^\lambda = S^\lambda \cap R$ for all $\lambda \in \Lambda$. To see this last statement, we use that $1_S \in S^0$, so that $R^\lambda = R^\lambda 1_S \in R^\lambda S^0 \subset S^\lambda$. Then $R^\lambda \subset S^\lambda \cap R$ follows. The other inclusion is immediate.

(b) We have seen in Theorem 6.5(b) that (B, κ_B) satisfies the IBF-principle. It follows that $R = \text{Span}_{\mathbb{Z}}\{\kappa_B(b_1, b_2) : b_i \in B\}$. Let now $\beta \in \text{IBF}_k(B)$ be a graded invariant bilinear form. Again by Theorem 6.5 there exists a unique $\varphi \in R^*$ such that $\beta = \varphi \circ \kappa_B$. We claim $\varphi(r) = 0$ for any $r \in R^\lambda$, $\lambda \neq 0$. Indeed, there exist finitely many $b_i \in B^{\mu_i}$ and $b'_i \in B^{\lambda-\mu_i}$ such that $r = \sum_i \kappa_B(b_i, b'_i)$. Hence $\varphi(r) = \sum_i \beta(b_i, b'_i) = 0$. That, conversely, every $\varphi \in (R^0)^*$ gives rise to a graded invariant bilinear form, is of course obvious. \square

Proposition 7.3 can be applied to multiloop algebras based on simple finite-dimensional Lie algebras. We recall their definition: \mathfrak{g} is a finite-dimensional simple Lie algebra over an algebraically closed field k of characteristic 0, and

$\sigma = (\sigma_1, \dots, \sigma_N)$ is a family of commuting automorphisms of \mathfrak{g} , which have finite orders m_1, \dots, m_N respectively. We fix a set of primitive m th roots of unity $\zeta_m \in k$ which are compatible in the sense that $\zeta_n^l = \zeta_n$, and put

$$R = k[t_1^{\pm 1}, \dots, t_N^{\pm 1}] \subset S = k[t_1^{\pm \frac{1}{m_1}}, \dots, t_N^{\pm \frac{1}{m_N}}].$$

The multiloop algebra $\mathcal{L} = \mathcal{L}(\mathfrak{g}, \sigma)$ associated to these data is the Lie algebra

$$\mathcal{L} = \bigoplus_{i_1, \dots, i_N \in \mathbb{Z}^N} \mathfrak{g}_{i_1, \dots, i_N} \otimes_k t_1^{\frac{i_1}{m_1}} \dots t_N^{\frac{i_N}{m_N}},$$

where $\mathfrak{g}_{i_1, \dots, i_N} = \{x \in \mathfrak{g} : \sigma_j(x) = \zeta_{m_j}^{i_j} x \text{ for all } j\}$. Since $\mathfrak{g}_{i_1, \dots, i_N} = \mathfrak{g}_{i_1+k_1, \dots, i_N+k_N}$ for $(k_1, \dots, k_N) \in m_1\mathbb{Z} \oplus \dots \oplus m_N\mathbb{Z}$, it is clear that \mathcal{L} is an R -Lie algebra. It is in fact an S/R -form of $\mathfrak{g} \otimes_k R$ (see [3, Theorem 3.6]).

To enter the grading into the picture, we let $\Lambda = \frac{1}{m_1}\mathbb{Z} \times \dots \times \frac{1}{m_N}\mathbb{Z}$. Then $\Lambda \simeq \mathbb{Z}^N$ and we have a natural Λ -grading on S and R (the reader will note that the homogeneous elements of R have degrees in $\mathbb{Z} \times \dots \times \mathbb{Z} \subset \Lambda$). The Lie algebra $\mathfrak{g} \otimes_k S$ is naturally Λ -graded and this makes \mathcal{L} also naturally into a Λ -graded Lie algebra. It is immediate from the definitions that \mathcal{L} is R -graded and a graded S/R -form of $\mathfrak{g} \otimes_k S$. Since in our situation $R^0 = k$, Proposition 7.3 yields the first part of the following.

Corollary 7.4. *Let \mathcal{L} be a multiloop Lie algebra based on a simple finite-dimensional Lie algebra over an algebraically closed field k of characteristic 0. Then, up to scalars in k , the \mathbb{Z}^N -graded Lie algebra \mathcal{L} has a unique graded invariant k -bilinear form β . It is given by*

$$\beta(x \otimes t_1^{\frac{j_1}{m_1}} \dots t_N^{\frac{j_N}{m_N}}, y \otimes t_1^{\frac{l_1}{m_1}} \dots t_N^{\frac{l_N}{m_N}}) = \kappa(x, y) \delta_{j_1+l_1, 0} \dots \delta_{j_N+l_N, 0},$$

where κ is the Killing form of \mathfrak{g} . The form β is nondegenerate. Every k -linear automorphism of \mathcal{L} is orthogonal with respect to β .

Proof. The nondegeneracy of κ implies that $\beta | \mathcal{L}^\lambda \times \mathcal{L}^{-\lambda}$ is a nondegenerate pairing for all λ , which in turn forces β to be nondegenerate.

Let now $f \in \text{Aut}_k(\mathcal{L})$. The map $\chi \mapsto f \circ \chi \circ f^{-1}$ is an automorphism of the centroid of \mathcal{L} . Since \mathcal{L} is central, f is α -semilinear for some k -linear automorphism α of R . By Proposition 6.1(b), we then know $\kappa_{\mathcal{L}} \circ (f \times f) = \alpha \circ \kappa_{\mathcal{L}}$ for $\kappa_{\mathcal{L}}$ the Killing form of the R -algebra \mathcal{L} . The form β is obtained by composing $\kappa_{\mathcal{L}}$ with the canonical projection $\epsilon : R \rightarrow R^0 = k$. It is therefore enough to show $\epsilon = \epsilon \circ \alpha$. But this is indeed the case: Every k -linear automorphism of R fixes R^0 pointwise and permutes the R^λ , $\lambda \neq 0$. To see this, we realize $\text{Aut}_k(R)$ as $\text{GL}_N(\mathbb{Z}) \ltimes (k^\times)^N$ in the natural fashion. □

This corollary is of interest for the construction of extended affine Lie algebras based on centerless Lie tori, which heavily depends on the existence of a graded

invariant nondegenerate bilinear form on a Lie torus. The reader is referred to [1, 20–22] for background material on extended affine Lie algebras and Lie tori. More precisely, we will deal here with Lie- \mathbb{Z}^N -tori.

Corollary 7.5. *Up to scalars, a centerless Lie- \mathbb{Z}^N -torus has a unique graded invariant nondegenerate bilinear form.*

Proof. By [19] a centerless Lie- \mathbb{Z}^N -torus is either finitely generated over its centroid or is a Lie torus with a root system of type A. Concerning the latter type, one knows from [4, 5, 31] that they are graded isomorphic to $\mathfrak{sl}_n(k_q)$ for a quantum torus k_q , and for these types of Lie algebras the claim follows from [22, 7.10]. If L is a centerless Lie- \mathbb{Z}^N -torus which is finitely generated over its centroid, then Theorem 3.3.1 of [2] says that L is graded-isomorphic to a multiloop algebra so that we can apply Corollary 7.4. □

Corollary 7.5 has been proven in [32, Theorem 7.1] for Lie tori graded by a torsion-free group Λ , using the structure theory of these types of Lie tori.

Acknowledgments

We thank Ottmar Loos for his many useful suggestions and comments. The first two authors gratefully acknowledge the hospitality of BIRS (Banff) and the Fields Institute (Toronto) where part of the research for this paper was carried out. E. Neher wishes to thank NSERC for partial support through a Discovery grant. A. Pianzola wishes to thank NSERC and CONICET for their continuous support. D. Prelat is supported by a Research Grant from Universidad CAECE. C. Sepp is supported by CONICET and Universidad CAECE.

References

- [1] B. Allison, S. Azam, S. Berman, Y. Gao and A. Pianzola, *Extended Affine Lie Algebras and Their Root Systems*, Memoirs of the American Mathematical Society, Vol. 126, No. 603 (American Mathematical Society, Providence, RI, 1997).
- [2] B. Allison, S. Berman, J. Faulkner and A. Pianzola, Multiloop realization of extended affine Lie algebras and Lie tori, *Trans. Amer. Math. Soc.* **361** (2009) 4807–4842.
- [3] B. Allison, S. Berman and A. Pianzola, Covering algebras. II. Isomorphism of loop algebras, *J. Reine Angew. Math.* **571** (2004) 39–71.
- [4] S. Berman, Y. Gao and Y. Krylyuk, Quantum tori and the structure of elliptic quasi-simple Lie algebras, *J. Funct. Anal.* **135** (1996) 339–389.
- [5] S. Berman, Y. Gao, Y. Krylyuk and E. Neher, The alternative torus and the structure of elliptic quasi-simple Lie algebras of type A_2 , *Trans. Amer. Math. Soc.* **347** (1995) 4315–4363.
- [6] R. Bix, Separable Jordan algebras over commutative rings, I, *J. Algebra* **57** (1979) 111–143.
- [7] ———, Separable alternative algebras over commutative rings, *J. Algebra* **92** (1985) 81–103.

- [8] N. Bourbaki, *Algèbre: Chapitres 1 à 3* (Masson, Paris, 1970).
- [9] ———, *Algèbre Commutative: Chapitres 1 à 4* (Masson, Paris, 1985).
- [10] F. DeMeyer and E. Ingraham, *Separable Algebras Over Commutative Rings*, Lecture Notes in Mathematics, Vol. 181 (Springer-Verlag, Berlin, 1971).
- [11] P. Gille and A. Pianzola, Galois cohomology and forms of algebras over Laurent polynomial rings, *Math. Ann.* **338**(2) (2007) 497–543.
- [12] B. Gross and G. Nebe, Globally maximal arithmetic groups, *J. Algebra* **272** (2004) 625–642.
- [13] M.-A. Knus and M. Ojanguren, *Théorie de la descente et algèbres d’Azumaya*, Lecture Notes in Mathematics, Vol. 389 (Springer-Verlag, Berlin, 1974), iv+163 pp.
- [14] O. Loos, *Jordan Pairs*, Lecture Notes in Mathematics, Vol. 460 (Springer-Verlag, Berlin, 1975), xvi+218 pp.
- [15] ———, Separable Jordan pairs over commutative rings, *Math. Ann.* **233** (1978) 137–144.
- [16] ———, Generically algebraic Jordan algebras over commutative rings, *J. Algebra* **297** (2006) 474–529.
- [17] O. Loos, H. Petersson and M. Racine, Inner derivations of alternative algebras over commutative rings, *Algebra Number Theory* **2** (2008) 927–968.
- [18] F. Montaner, A. Stolin and E. Zelmanov, Classification of Lie bialgebras over current algebras, *Selecta Math. (N.S.)* **16** (2010) 935–962.
- [19] E. Neher, Lie tori, *C. R. Math. Acad. Sci. Soc. R. Can.* **26** (2004) 84–89.
- [20] ———, Extended affine Lie algebras, *C. R. Math. Acad. Sci. Soc. R. Can.* **26** (2004) 90–96.
- [21] ———, Extended affine Lie algebras — an introduction to their structure theory, in *Geometric Representation Theory and Extended Affine Lie Algebras*, Fields Institute Communications, Vol. 59 (American Mathematical Society, Providence, RI, 2011), pp. 107–167.
- [22] ———, Extended affine Lie algebras and other generalizations of affine Lie algebras — a survey, in *Developments and Trends in Infinite-Dimensional Lie Theory*, Progress in Mathematics, Vol. 288 (Birkhäuser Boston Inc., Boston, MA, 2011), pp. 53–126.
- [23] E. Neher and A. Pianzola, Étale descent of derivations, *Transformations Groups* **18**(4) (2013) 1189–1205.
- [24] H. Petersson, Composition algebras over algebraic curves of genus zero, *Trans. Amer. Math. Soc.* **337** (1993) 473–493.
- [25] A. Pianzola, Derivations of certain algebras defined by étale, descent, *Math. Z.* **264** (2010) 485–495.
- [26] ———, Vanishing of H^1 for Dedekind rings and applications to loop algebras, *C. R. Math. Acad. Sci. Paris* **340** (2005) 633–638.
- [27] A. Pianzola, D. Prelat and C. Sepp, Variations on themes of C. Kassel and R. Wilson, in preparation.
- [28] A. Grothendieck, *Séminaire de Géométrie Algébrique de l’I.H.E.S., Revêtements étales et groupe fondamental*, Lecture Notes in Mathematics, Vol. 224 (Springer, 1971).
- [29] T. A. Springer and R. Steinberg, Conjugacy classes, in *Seminar on Algebraic Groups and Related Finite Groups*, Lecture Notes in Mathematics, Vol. 131 (Springer-Verlag, 1970), pp. 167–266.
- [30] W. C. Waterhouse, *Introduction to Affine Group Schemes*, Graduate Texts in Mathematics, Vol. 66 (Springer, New York, 1979).

- [31] Y. Yoshii, Coordinate algebras of extended affine Lie algebras of type A_1 , *J. Algebra* **234** (2000) 128–168.
- [32] ———, Lie tori — a simple characterization of extended affine Lie algebras, *Publ. Res. Inst. Math. Sci.* **42** (2006) 739–762.
- [33] P. Zusmanovich, The second homology group of current Lie algebras, *Astérisque* **226** (1994) 435–452.